

CYBR1 mouse mAb


Catalog No YP-mAb-12238 Isotype IgG Reactivity Human; Mouse;Rat Applications WB Gene Name CYBRD1 DCYTB FRRS3 Protein Name CYBR1 Immunogen Synthesized peptide derived from human CYBR1 AA ranges Specificity This antibody detects endogenous levels of CYBR1 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.029 Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90%	41-91
ReactivityHuman; Mouse;RatApplicationsWBGene NameCYBRD1 DCYTB FRRS3Protein NameCYBR1ImmunogenSynthesized peptide derived from human CYBR1 AA range: SpecificitySpecificityThis antibody detects endogenous levels of CYBR1 at HumanFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02%SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-1:2000Concentration1 mg/ml	41-91
Applications Gene Name CYBRD1 DCYTB FRRS3 Protein Name CYBR1 Immunogen Synthesized peptide derived from human CYBR1 AA range: Specificity This antibody detects endogenous levels of CYBR1 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.029 Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml	41-91
Gene Name CYBRD1 DCYTB FRRS3 Protein Name CYBR1 Immunogen Synthesized peptide derived from human CYBR1 AA range: Specificity This antibody detects endogenous levels of CYBR1 at Human Experimental Ex	41-91
Protein Name CYBR1 Immunogen Synthesized peptide derived from human CYBR1 AA range: This antibody detects endogenous levels of CYBR1 at Human Experimental Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.029 Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml	41-91
ImmunogenSynthesized peptide derived from human CYBR1 AA range:SpecificityThis antibody detects endogenous levels of CYBR1 at Human Equipment In the Indian Equipment In t	41-91
SpecificityThis antibody detects endogenous levels of CYBR1 at HumanFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02%SourceMonoclonal, Mouse, lgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-1:2000Concentration1 mg/ml	41-91
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.029 Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml	
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml	ın/Mouse/Rat
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml	် sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml	
Concentration 1 mg/ml	
Purity ≥90%	
Storage Stability -20°C/1 year	
Synonyms	
Observed Band	
Cell Pathway Cell membrane ; Multi-pass membrane protein . Apical cell r Multi-pass membrane protein . Localized at the brush border	nembrane ; of duodenal cells
Tissue Specificity Present in erythrocyte membranes (at protein level). Also ex epithelium.	pressed in respiratory
cofactor:Binds 2 heme groups non-covalently.,disease:Defebe a cause of primary hereditary hemochromatosis (HFE). He disorder characterized by iron accumulation in parenchymal accumulation usually results in tissue damage and causes of diabetes mellitus, arthropathy, cardiomyopathy, endocrine a increased risk of hepatocellular carcinoma.,function:Ferric-creduces Fe(3+) to Fe(2+). Present at the brush border of due where it probably reduces dietary Fe(3+) thereby facilitating mucosal cells. Uses ascorbate as electron donor. May be in ascorbate recycling in erythrocyte membranes. May also act airway epithelial cells.,induction:By iron deficiency (at protein level).,similarity:Contains 1 cytochrom	FE is an iron-loading cells. Iron rrhosis of the liver, onormalities and an nelate reductase that odenal enterocytes its transport into the
Background This gene is a member of the cytochrome b(561) family that iron-regulated protein. It highly expressed in the duodenal by membrane. It has ferric reductase activity and is believed to	as a ferrireductase in



UpingBio technology Co.,Ltd



role in dietary iron absorption. [provided by RefSeq, Jul 2008],



Products Images

