






ABR mouse mAb

them.,similarity:Contains 1 C2 domain.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 Rho-GAP domain.,tissue specificity:Highly enriched in the brain. Much weaker expression in heart, lung and muscle., This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating protein domain, a domain found in members of the Rho family of GTP-binding proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively		
Reactivity Human; Mouse Applications WB Gene Name ABR Protein Name ABR Immunogen Synthesized peptide derived from human ABR AA range: 393-443 Specificity This antibody detects endogenous levels of ABR at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell projection, dendritic spine. Cell projection, axon. Cell junction, synapse. Tissue Specificity Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Function function:GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them, similarity Contains 1 C2 domain, similarity-Contains 1 Rho-GAP domain, lissue specificity. Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Background This gene en	Catalog No	YP-mAb-12298
Applications Gene Name ABR Protein Name ABR Immunogen Synthesized peptide derived from human ABR AA range: 393-443 Specificity This antibody detects endogenous levels of ABR at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell projection, dendritic spine . Cell projection, axon . Cell junction, synapse . Tissue Specificity Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Function function:GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them, similarity:Contains 1 PH domain, similarity:Contains 1 DH (DBL-nomology) domain. similarity:Contains 1 PH domain, similarity:Contains 1 Rho-GAP domain, lissue specificity:Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Background This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating protein. Sunctional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene, [provided by RefSeq, 9].	Isotype	IgG
Gene Name ABR Protein Name ABR Immunogen Synthesized peptide derived from human ABR AA range: 393-443 Specificity This antibody detects endogenous levels of ABR at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell projection, dendritic spine . Cell projection, axon . Cell junction, synapse . Tissue Specificity Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Function function:GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them, similarity:Contains 1 DH (DBL-nomology) domain, similarity:Contains 1 DH domain, similarity:Contains 1 Rho-GAP domain, ilssue specificity:Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Background This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating protein domain, adomain found in members of the Rho family of GTP-binding proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene, [provided by RefSeq, provided by RefS	Reactivity	Human; Mouse
Protein Name ABR	Applications	WB
Immunogen Synthesized peptide derived from human ABR AA range: 393-443 Specificity This antibody detects endogenous levels of ABR at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell projection, dendritic spine . Cell projection, axon . Cell junction, synapse . Tissue Specificity Highly enriched in the brain. Much weaker expression in heart, lung and muscle . Function function:GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating 1 DH (DBL-homology) domain, similarity:Contains 1 C2 domain, similarity:Contains 1 DH (DBL-homology) domain, issue specificity:Highly enriched in the brain. Much weaker expression in heart, lung and muscle . Background This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating protein formain, a domain found in members of the Rho family of GTP-	Gene Name	ABR
Specificity This antibody detects endogenous levels of ABR at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell projection, dendritic spine. Cell projection, axon. Cell junction, synapse. Tissue Specificity Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Function function:GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them, similarity:Contains 1 C2 domain, similarity:Contains 1 DH (DBL-homology) domain, similarity:Contains 1 C2 domain, similarity:Contains 1 DH (DBL-homology) domain, similarity:Contains 1 PH domain, similarity:Contains 1 Rho-GAP Background This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene encotains a GTPase-activating protein domain, a domain found in members of the Rho family of GTP-binding protein semicolaed admain found in members of the Rho family of GTP-binding protein omain, a domain found in members of the Rho family of GTP-binding protein domain, a domain found in members of the Rho family of GTP-binding protein domain, a domain found in members of the Rho family of GTP-binding protein formain, a domain found in members of the Rho family of GTP-binding protein domain, a domain found in members of the Rho family of GTP-binding protein domain, a domain found in members of the Rho family of gtp. provided by RefSeq. provided by RefSeq.	Protein Name	ABR
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell projection, dendritic spine . Cell projection, axon . Cell junction, synapse . Tissue Specificity Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Function function:GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them, similarity. Contains 1 DH (DBL-homology) domain, similarity. Contains 1 PH domain. similarity. Contains 1 Ph domain. similarity. Contains 1 Rho-GAP domain, similarity. Contains 1 PH domain. similarity. Contains 1 Rho-GAP domain, tissue specificity:Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Background This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating protein domain, a domain found in members of the Rho family of GTP-binding proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene. [provided by RefSeq,	Immunogen	Synthesized peptide derived from human ABR AA range: 393-443
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell projection, dendritic spine . Cell projection, axon . Cell junction, synapse . Tissue Specificity Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Function function:GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them.,similarity:Contains 1 C2 domain.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 Rho-GAP domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 Rho-GAP domain.,similarity:Contains 1 Rho-GAP domain. and muscle. Background This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating protein domain, a domain found in members of the Rho family of GTP-binding proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene. [provided by RefSeq,	Specificity	This antibody detects endogenous levels of ABR at Human/Mouse
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell projection, dendritic spine . Cell projection, axon . Cell junction, synapse . Tissue Specificity Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Function function:GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 Rho-GAP domain.,issue specificity:Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Background This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcripty variants have been reported for this gene. [provided by RefSeq.]	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell projection, dendritic spine . Cell projection, axon . Cell junction, synapse . Tissue Specificity Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Function function:GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them., similarity:Contains 1 DH (DBL-homology) domain., similarity:Contains 1 PH domain., similarity:Contains 1 Rho-GAP domain., isimilarity:Contains 1 PH domain., similarity:Contains 1 Rho-GAP domain., isimilarity:Contains 2 Rho-GAP domain., isimilar	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell projection, dendritic spine . Cell projection, axon . Cell junction, synapse . Tissue Specificity Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Function function:GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them.,similarity:Contains 1 C2 domain.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 Rho-GAP domain.,tissue specificity:Highly enriched in the brain. Much weaker expression in heart, lung and muscle., Background This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene. [provided by RefSeq.]	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell projection, dendritic spine . Cell projection, axon . Cell junction, synapse . Tissue Specificity Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Function function:GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them.,similarity:Contains 1 C2 domain.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 Rho-GAP domain.,tissue specificity:Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Background This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene. [provided by RefSeq,	Dilution	WB 1:500-1:2000
Synonyms Observed Band Cell Pathway Cell projection, dendritic spine. Cell projection, axon. Cell junction, synapse. Tissue Specificity Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Function function:GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them., similarity:Contains 1 C2 domain., similarity:Contains 1 DH (DBL-homology) domain., similarity:Contains 1 PH domain., similarity:Contains 1 Rho-GAP domain., tissue specificity:Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Background This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating protein domain, a domain found in members of the Rho family of GTP-binding proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene. [provided by RefSeq.]	Concentration	1 mg/ml
Observed Band Cell Pathway Cell projection, dendritic spine. Cell projection, axon. Cell junction, synapse. Tissue Specificity Highly enriched in the brain. Much weaker expression in heart, lung and muscle. Function function:GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them.,similarity:Contains 1 C2 domain.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 Rho-GAP domain.,tissue specificity:Highly enriched in the brain. Much weaker expression in heart, lung and muscle., Background This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene. [provided by RefSeq,	Purity	≥90%
Observed Band Cell Pathway Cell projection, dendritic spine. Cell projection, axon. Cell junction, synapse. Tissue Specificity Highly enriched in the brain. Much weaker expression in heart, lung and muscle. function: GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them., similarity: Contains 1 C2 domain., similarity: Contains 1 DH (DBL-homology) domain., similarity: Contains 1 PH domain., similarity: Contains 1 Rho-GAP domain., tissue specificity: Highly enriched in the brain. Much weaker expression in heart, lung and muscle., Background This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating protein domain, a domain found in members of the Rho family of GTP-binding proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene. [provided by RefSeq,	Storage Stability	-20°C/1 year
Cell Pathway Cell projection, dendritic spine. Cell projection, axon. Cell junction, synapse. Highly enriched in the brain. Much weaker expression in heart, lung and muscle. function:GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them.,similarity:Contains 1 C2 domain.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 Rho-GAP domain.,tissue specificity:Highly enriched in the brain. Much weaker expression in heart, lung and muscle., Background This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating protein domain, a domain found in members of the Rho family of GTP-binding proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene. [provided by RefSeq,	Synonyms	
Tissue Specificity Highly enriched in the brain. Much weaker expression in heart, lung and muscle. function function:GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them.,similarity:Contains 1 C2 domain.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 Rho-GAP domain.,tissue specificity:Highly enriched in the brain. Much weaker expression in heart, lung and muscle., This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating protein domain, a domain found in members of the Rho family of GTP-binding proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene. [provided by RefSeq,	Observed Band	
function: GTPase-activating protein for RAC and CDC42. Promotes the exchange of RAC or CDC42-bound GDP by GTP, thereby activating them., similarity: Contains 1 C2 domain., similarity: Contains 1 DH (DBL-homology) domain., similarity: Contains 1 PH domain., similarity: Contains 1 Rho-GAP domain., tissue specificity: Highly enriched in the brain. Much weaker expression in heart, lung and muscle., Background This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating protein domain, a domain found in members of the Rho family of GTP-binding proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene. [provided by RefSeq,	Cell Pathway	Cell projection, dendritic spine . Cell projection, axon . Cell junction, synapse .
them.,similarity:Contains 1 C2 domain.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 Rho-GAP domain.,tissue specificity:Highly enriched in the brain. Much weaker expression in heart, lung and muscle., This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating protein domain, a domain found in members of the Rho family of GTP-binding proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene. [provided by RefSeq,	Tissue Specificity	Highly enriched in the brain. Much weaker expression in heart, lung and muscle.
breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating protein domain, a domain found in members of the Rho family of GTP-binding proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene. [provided by RefSeq,	Function	domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 Rho-GAP domain.,tissue specificity:Highly enriched in the brain. Much weaker expression in
	Background	breakpoint cluster region gene located on chromosome 22. The protein encoded by this gene contains a GTPase-activating protein domain, a domain found in members of the Rho family of GTP-binding proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene. [provided by RefSeq,



UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using ABR mouse mAb