



NF-M Monoclonal Antibody

Catalog No	YP-mAb-12766
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	NEFM
Protein Name	Neurofilament medium polypeptide
Immunogen	The antiserum was produced against synthesized peptide derived from human NF-M. AA range:542-591
Specificity	NF-M Monoclonal Antibody detects endogenous levels of NF-M protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NEFM; NEF3; NFM; Neurofilament medium polypeptide; NF-M; 160 kDa neurofilament protein; Neurofilament 3; Neurofilament triplet M protein
Observed Band	110kD
Cell Pathway	Cytoplasm, cytoskeleton . Cell projection, axon .
Tissue Specificity	Brain,Brain cortex,Fetal brain cortex,
Function	function:Neurofilaments usually contain three intermediate filament proteins: L, M, and H which are involved in the maintenance of neuronal caliber.,PTM:Phosphorylation seems to play a major role in the functioning of the larger neurofilament polypeptides (NF-M and NF-H), the levels of phosphorylation being altered developmentally and coincident with a change in the neurofilament function.,PTM:There are a number of repeats of the tripeptide K-S-P, NFM is phosphorylated on a number of the serines in this motif. It is thought that phosphorylation of NFM results in the formation of interfilament cross bridges that are important in the maintenance of axonal caliber.,similarity:Belongs to the intermediate filament family.,
Background	neurofilament, medium polypeptide(NEFM) Homo sapiens Neurofilaments are type IV intermediate filament heteropolymers composed of light, medium, and heavy chains. Neurofilaments comprise the axoskeleton and functionally maintain neuronal caliber. They may also play a role in intracellular transport to axons and



dendrites. This gene encodes the medium neurofilament protein. This protein is commonly used as a biomarker of neuronal damage. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Oct 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using NF-M Monoclonal Antibody