



## NTR2 Monoclonal Antibody

Catalog No         YP-mAb-12782           Isotype         IgG           Reactivity         Human;Mouse;Rat           Applications         WB           Gene Name         NTSR2           Protein Name         Neurotensin receptor type 2           Immunogen         The antiserum was produced against synthesized peptide derived from human NTR2. AA range:151-200           Specificity         NTR2 Monoclonal Antibody detects endogenous levels of NTR2 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         NTSR2, Neurotensin receptor type 2; NT-R-2; NTR2; Levocabastine-sensitive neurotensin receptor           Observed Band         45kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Expressed in prostate (at protein level).           Function         function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that		
Reactivity Human;Mouse;Rat  Applications WB  Gene Name NTSR2  Protein Name Neurotensin receptor type 2  Immunogen The antiserum was produced against synthesized peptide derived from human NTR2. AA range:151-200  Specificity NTR2 Monoclonal Antibody detects endogenous levels of NTR2 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms NTSR2; Neurotensin receptor type 2; NT-R-2; NTR2; Levocabastine-sensitive neurotensin receptor  Observed Band 45kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity Expressed in prostate (at protein level).  Function function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphaldylinositol-calcium second messenger system, online information Neurotensin receptor entry, similarity:Belongs to the G-protein coupled receptor 1 family, and activate a phosphaldylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor inamily that activate a phosphaldylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor inamily that activate a phosphaldylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor inamily that receptor algorables and several other lagorables are recognizes, with his receptor. However, unlike NT1 receptor, this gene recognizes, with his receptor inable of protein-coupled freceptor and be of physiological importance and finity, Incocabastine, a histamine HT receptor and postpart described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with historices of may be of physiological importance and that a natural sun	Catalog No	YP-mAb-12782
Applications WB  Gene Name NTSR2  Protein Name Neurotensin receptor type 2  Immunogen The antiserum was produced against synthesized peptide derived from human NTR2. AA range:151-200  Specificity NTR2 Monoclonal Antibody detects endogenous levels of NTR2 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity 290%  Storage Stability -20°C/1 year  Synonyms NTSR2; Neurotensin receptor type 2; NT-R-2; NTR2; Levocabastine-sensitive neurotensin receptor  Observed Band 45kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity Expressed in prostate (at protein level).  Function function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system., online information:Neurotensin receptor entry, similarity, Belongs to the G-protein coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor pridusing in affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural	Isotype	IgG
Gene Name         NTSR2           Protein Name         Neurotensin receptor type 2           Immunogen         The antiserum was produced against synthesized peptide derived from human NTR2. An range:151-200           Specificity         NTR2 Monoclonal Antibody detects endogenous levels of NTR2 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse,lgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         NTSR2; Neurotensin receptor type 2; NT-R-2; NTR2; Levocabastine-sensitive neurotensin receptor           Observed Band         45kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Expressed in prostate (at protein level).           Function         function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system. Jonilien information:Neurotensin receptor entry, similarity:Belongs to the G-protein coupled receptor 1 family.           Background         The protein encoded by this gene belongs to the G	Reactivity	Human;Mouse;Rat
Immunogen	Applications	WB
Immunogen         The antiserum was produced against synthesized peptide derived from human NTR2. AA range:151-200           Specificity         NTR2 Monoclonal Antibody detects endogenous levels of NTR2 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         NTSR2; Neurotensin receptor type 2; NT-R-2; NTR2; Levocabastine-sensitive neurotensin receptor           Observed Band         45kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Expressed in prostate (at protein level).           Function         function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system. Joining in protein in a country similarity:Belongs to the G-protein coupled receptor 1 family.           Background         The protein encoded by this gene belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurot	Gene Name	NTSR2
NTR2. AA range:151-200  Specificity NTR2 Monoclonal Antibody detects endogenous levels of NTR2 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms NTSR2; Neurotensin receptor type 2; NT-R-2; NTR2; Levocabastine-sensitive neurotensin receptor  Observed Band 45kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Expressed in prostate (at protein level).  Function function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system., online information:Neurotensin receptor entry, similarity:Belongs to the G-protein coupled receptor 1 family.  The protein encoded by this gene belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor linds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor and that a natural	Protein Name	Neurotensin receptor type 2
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms NTSR2; Neurotensin receptor type 2; NT-R-2; NTR2; Levocabastine-sensitive neurotensin receptor  Observed Band 45kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity Expressed in prostate (at protein level).  Function function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system, online information:Neurotensin receptor entry,similarity:Belongs to the G-protein coupled receptor 1 family.  Background The protein encoded by this gene belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural	Immunogen	
Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         NTSR2; Neurotensin receptor type 2; NT-R-2; NTR2; Levocabastine-sensitive neurotensin receptor           Observed Band         45kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Expressed in prostate (at protein level).           Function         function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system. Jonline information:Neurotensin receptor entry, similarity:Belongs to the G-protein coupled receptor 1 family.           Background         The protein encoded by this gene belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin neurotensin as well as several other ligands already described for neurotensin affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural </th <th>Specificity</th> <th>NTR2 Monoclonal Antibody detects endogenous levels of NTR2 protein.</th>	Specificity	NTR2 Monoclonal Antibody detects endogenous levels of NTR2 protein.
Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-1:2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  NTSR2; Neurotensin receptor type 2; NT-R-2; NTR2; Levocabastine-sensitive neurotensin receptor  Observed Band  45kD  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Expressed in prostate (at protein level).  Function  function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system, online information:Neurotensin receptor entry, similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  The protein encoded by this gene belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms NTSR2; Neurotensin receptor type 2; NT-R-2; NTR2; Levocabastine-sensitive neurotensin receptor  Observed Band 45kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity Expressed in prostate (at protein level).  Function function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system., online information:Neurotensin receptor entry, similarity:Belongs to the G-protein coupled receptor 1 family.,  Background The protein encoded by this gene belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural	Source	Monoclonal, Mouse,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       NTSR2; Neurotensin receptor type 2; NT-R-2; NTR2; Levocabastine-sensitive neurotensin receptor         Observed Band       45kD         Cell Pathway       Cell membrane; Multi-pass membrane protein.         Tissue Specificity       Expressed in prostate (at protein level).         Function       function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system, online information:Neurotensin receptor entry,similarity:Belongs to the G-protein coupled receptor 1 family.         Background       The protein encoded by this gene belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity bindings lites in brain. These activities suggest that this receptor may be of physiological importance and that a natural	Purification	·
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms NTSR2; Neurotensin receptor type 2; NT-R-2; NTR2; Levocabastine-sensitive neurotensin receptor  Observed Band 45kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity Expressed in prostate (at protein level).  Function function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system., online information:Neurotensin receptor entry, similarity:Belongs to the G-protein coupled receptor 1 family.  Background The protein encoded by this gene belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural	Dilution	WB 1:500-1:2000
Storage Stability  -20°C/1 year  NTSR2; Neurotensin receptor type 2; NT-R-2; NTR2; Levocabastine-sensitive neurotensin receptor  Observed Band  45kD  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Expressed in prostate (at protein level).  Function  function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system., online information:Neurotensin receptor entry, similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  The protein encoded by this gene belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural	Concentration	1 mg/ml
Synonyms  NTSR2; Neurotensin receptor type 2; NT-R-2; NTR2; Levocabastine-sensitive neurotensin receptor  Observed Band  45kD  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Expressed in prostate (at protein level).  Function  function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system., online information:Neurotensin receptor entry, similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  The protein encoded by this gene belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural	Purity	≥90%
Deserved Band  45kD  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Expressed in prostate (at protein level).  Function  function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system.,online information:Neurotensin receptor entry,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  The protein encoded by this gene belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural	Storage Stability	-20°C/1 year
Cell Pathway  Cell membrane; Multi-pass membrane protein.  Expressed in prostate (at protein level).  Function  function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system.,online information:Neurotensin receptor entry,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  The protein encoded by this gene belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural	Synonyms	· · · · · · · · · · · · · · · · · · ·
Tissue Specificity  Expressed in prostate (at protein level).  function  function:Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system.,online information:Neurotensin receptor entry,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  The protein encoded by this gene belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural	Observed Band	45kD
function: Receptor for the tridecapeptide neurotensin. It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system., online information: Neurotensin receptor entry, similarity: Belongs to the G-protein coupled receptor 1 family.,  Background  The protein encoded by this gene belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural	Cell Pathway	Cell membrane; Multi-pass membrane protein.
proteins that activate a phosphatidylinositol-calcium second messenger system.,online information:Neurotensin receptor entry,similarity:Belongs to the G-protein coupled receptor 1 family.,  The protein encoded by this gene belongs to the G protein-coupled receptor family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural	Tissue Specificity	Expressed in prostate (at protein level).
family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural	Function	proteins that activate a phosphatidylinositol-calcium second messenger system.,online information:Neurotensin receptor entry,similarity:Belongs to the
	Background	family that activate a phosphatidylinositol-calcium second messenger system. Binding and pharmacological studies demonstrate that this receptor binds neurotensin as well as several other ligands already described for neurotensin NT1 receptor. However, unlike NT1 receptor, this gene recognizes, with high affinity, levocabastine, a histamine H1 receptor antagonist previously shown to compete with neurotensin for low-affinity binding sites in brain. These activities suggest that this receptor may be of physiological importance and that a natural



## UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





-- 85

Western Blot analysis of various cells using NTR2 Monoclonal Antibody

NTR2

-- 48

-- 34

-- 26

-- 19

(kD)