



Histamine H4 Receptor Monoclonal Antibody

Catalog No	YP-mAb-13362
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	HRH4
Protein Name	Histamine H4 receptor
Immunogen	The antiserum was produced against synthesized peptide derived from human HRH4. AA range:221-270
Specificity	Histamine H4 Receptor Monoclonal Antibody detects endogenous levels of Histamine H4 Receptor protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HRH4; GPCR105; Histamine H4 receptor; H4R; HH4R; AXOR35; G-protein coupled receptor 105; GPRv53; Pfi-013; SP9144
Observed Band	45kD
Cell Pathway	Cell membrane; Multi-pass membrane protein.
Tissue Specificity	Expressed primarily in the bone marrow and eosinophils. Shows preferential distribution in cells of immunological relevance such as T-cells, dendritic cells, monocytes, mast cells, neutrophils. Also expressed in a wide variety of peripheral tissues, including the heart, kidney, liver, lung, pancreas, skeletal muscle, prostate, small intestine, spleen, testis, colon, fetal liver and lymph node.
Function	function:The H4 subclass of histamine receptors could mediate the histamine signals in peripheral tissues. Displays a significant level of constitutive activity (spontaneous activity in the absence of agonist).,induction:Expression is either up-regulated or down-regulated upon activation of the lymphoid tissues and this regulation may depend on the presence of IL-10 or IL-13.,miscellaneous:Does not bind diphenhydramine, loratadine, ranitidine, cimetidine and chlorpheniramine. Shows modest affinity for dimaprit, impromidine, clobenpropit, thioperamide, burimamide clozapine, immepip and imetit. The order of inhibitory activity was imetit > clobenpropit > burimamide > thioperamide. Clobenpropit behaves as a partial agonist, dimaprit and impromidine show some agonist activity while clozapine behaves as a full agonist. Thioperamide shows inverse agonism



(enhances cAMP activity). The order of

Background

Histamine is a ubiquitous messenger molecule released from mast cells, enterochromaffin-like cells, and neurons. Its various actions are mediated by a family of histamine receptors, which are a subset of the G-protein coupled receptor superfamily. This gene encodes a histamine receptor that is predominantly expressed in haematopoietic cells. The protein is thought to play a role in inflammation and allergy responses. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Histamine H4 Receptor Monoclonal Antibody