





Olfactory receptor 10G4 Monoclonal Antibody

Catalog No YP-mAb-13441 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name OR10G4 Protein Name Olfactory receptor 10G4 Immunogen The antiserum was produced against synthesized peptide derived from human OR10G4. AA range:230-279 Specificity Olfactory receptor 10G4 Monoclonal Antibody detects endogenous levels of Olfactory receptor 10G4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., arising from single coding-e		
Reactivity Human;Rat;Mouse; Applications WB Gene Name OR10G4 Protein Name Olfactory receptor 10G4 Immunogen The antiserum was produced against synthesized peptide derived from human OR10G4. AA range:230-279 Specificity Olfactory receptor 10G4 Monoclonal Antibody detects endogenous levels of Olfactory receptor 10G4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family. Difactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are a members of a large family of G-protein-coupled receptors of a large family of G-protein-coupled receptors or proteins are nembers of a large family of G-protein-coupled receptors or proteins are nembers of a large family of G-protein-coupled receptors or a responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by proteins are norman is independent of other organisms. [provided by proteins and proteins for this organism is independent of other organisms.] provided by	Catalog No	YP-mAb-13441
Applications Gene Name OR10G4 Protein Name Olfactory receptor 10G4 Immunogen The antiserum was produced against synthesized peptide derived from human OR10G4. AA range:230-279 Specificity Olfactory receptor 10G4 Monoclonal Antibody detects endogenous levels of Olfactory receptor 10G4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor septes and proteins for this organism is independent of other organisms. [provided by proteins for this organism is independent of other organisms. [provided by	Isotype	IgG
Gene Name OR10G4 Protein Name Olfactory receptor 10G4 Immunogen The antiserum was produced against synthesized peptide derived from human OR10G4. AA range:230-279 Specificity Olfactory receptor 10G4 Monoclonal Antibody detects endogenous levels of Olfactory receptor 10G4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organism is independent of other organisms, provided by	Reactivity	Human;Rat;Mouse;
Protein Name Olfactory receptor 10G4 Immunogen The antiserum was produced against synthesized peptide derived from human OR10G4. AA range:230-279 Specificity Olfactory receptor 10G4 Monoclonal Antibody detects endogenous levels of Olfactory receptor 10G4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organism is independent of other organisms, provided by	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human OR10G4. AA range:230-279 Specificity Olfactory receptor 10G4 Monoclonal Antibody detects endogenous levels of Olfactory receptor 10G4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor green arriving from single coding-exon genes. Olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors smell. The olfactory receptors and hormone receptors and are responsible for the recognition and G protein-mediated transduction of oddrant signals. The olfactory receptor g	Gene Name	OR10G4
Specificity Olfactory receptor 10G4 Monoclonal Antibody detects endogenous levels of Olfactory receptor 10G4 Monoclonal Antibody detects endogenous levels of Olfactory receptor 10G4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor gene family is the largest in the genome. The nomenclature assigned	Protein Name	Olfactory receptor 10G4
Tissue Specificity Function Olfactory receptor 10G4 protein. Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function: Odorant receptor ., similarity: Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors capture a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Immunogen	
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor ,,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms, [provided by	Specificity	Olfactory receptor 10G4 Monoclonal Antibody detects endogenous levels of Olfactory receptor 10G4 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function: Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Dilution	WB 1:500-1:2000
Synonyms OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Concentration	1 mg/ml
Synonyms OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Purity	≥90%
Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Synonyms	OR10G4; Olfactory receptor 10G4; Olfactory receptor OR11-278
Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Observed Band	35kD
Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Cell Pathway	Cell membrane; Multi-pass membrane protein.
Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Tissue Specificity	
neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Function	
	Background	neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by



UpingBio technology Co.,Ltd







matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

