







Olfactory receptor 8G2 Monoclonal Antibody

	VP AL 40000
Catalog No	YP-mAb-13632
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	OR8G2
Protein Name	Olfactory receptor 8G2
Immunogen	The antiserum was produced against synthesized peptide derived from human OR8G2. AA range:237-286
Specificity	Olfactory receptor 8G2 Monoclonal Antibody detects endogenous levels of Olfactory receptor 8G2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,lgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	OR8G2; OR8G2P; OR8G4; Olfactory receptor 8G2; Olfactory receptor 8G4; Olfactory receptor OR11-297; Olfactory receptor TPCR120
Observed Band	33kD
Cell Pathway	plasma membrane,integral component of membrane,
Tissue Specificity	Pooled,Testicle,Testis,
Function	function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
Background	olfactory receptor family 8 subfamily G member 2(OR8G2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],



UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





-- 85

Western Blot analysis of various cells using Olfactory receptor 8G2 Monoclonal Antibody

OR8G2

-- 48

-- 26

(kD)