





## T2R43 Monoclonal Antibody

Catalog No	YP-mAb-13684
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	TAS2R43
Protein Name	Taste receptor type 2 member 43
Immunogen	The antiserum was produced against synthesized peptide derived from human TAS2R43. AA range:124-173
Specificity	T2R43 Monoclonal Antibody detects endogenous levels of T2R43 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TAS2R43; Taste receptor type 2 member 43; T2R43; Taste receptor type 2 member 52; T2R52
Observed Band	35kD
Cell Pathway	Membrane ; Multi-pass membrane protein . Cell projection, cilium membrane . In airway epithelial cells, localizes to motile cilia.
Tissue Specificity	Expressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells. Expressed in airway epithelia.
Function	function:Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5 (By similarity). Activated by the sulfonyl amide sweeteners saccharin and acesulfame K.,miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and exclusively in gustducin-positive cells.,
Background	TAS2R43 belongs to the large TAS2R receptor family. TAS2Rs are expressed on the surface of taste receptor cells and mediate the perception of bitterness through a G protein-coupled second messenger pathway (Conte et al., 2002



## UpingBio technology Co.,Ltd





[PubMed 12584440]). For further information on TAS2Rs, see MIM 604791.[supplied by OMIM, Mar 2009],

matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

TAS2R43

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Monoclonal Antibody

Western Blot analysis of various cells using T2R43





--250 --150

--100

--75

--50

--37

--25

--20

--15

(kd)

JK

Thank you for your recent purchase

Focus on your research Service for life science