





TAAR3 Monoclonal Antibody

YP-mAb-13691
IgG
Human;Rat;Mouse;
WB
TAAR3
Putative trace amine-associated receptor 3
The antiserum was produced against synthesized peptide derived from human TAAR3. AA range:206-255
TAAR3 Monoclonal Antibody detects endogenous levels of TAAR3 protein.
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-1:2000
1 mg/ml
≥90%
-20°C/1 year
TAAR3; GPR57; Putative trace amine-associated receptor 3; TaR-3; Trace amine receptor 3; G-protein coupled receptor 57
39kD
Cell membrane ; Multi-pass membrane protein .
Not expressed in the pons, thalamus, globus pallidus, caudate, putamen or cerebellum.
caution:Seems to be a non functional due to its divergence in positions 20 to 45 (due to a frameshift) but does not seem to be the product of a pseudogene.,function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Not expressed in the pons, thalamus, globus pallidus, caudate, putamen or cerebellum.,
caution:Seems to be a non functional due to its divergence in positions 20 to 45 (due to a frameshift) but does not seem to be the product of a pseudogene.,function:Orphan receptor.,similarity:Belongs to the G-protein
coupled receptor 1 family.,tissue specificity:Not expressed in the pons, thalamus, globus pallidus, caudate, putamen or cerebellum.,



UpingBio technology Co.,Ltd








Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using TAAR3 Monoclonal Antibody