




Fgl2 Monoclonal Antibody

Catalog No	YP-mAb-13927
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	FGL2
Protein Name	Fibroleukin
Immunogen	The antiserum was produced against synthesized peptide derived from human Fgl2. AA range:38-87
Specificity	Fgl2 Monoclonal Antibody detects endogenous levels of Fgl2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	FGL2; Fibroleukin; Fibrinogen-like protein 2; pT49
Observed Band	55kD
Cell Pathway	Secreted.
Tissue Specificity	Constitutively expressed in cytotoxic T-cells.
Function	function:May play a role in physiologic lymphocyte functions at mucosal sites.,similarity:Contains 1 fibrinogen C-terminal domain.,subunit:Homotetramer; disulfide-linked.,tissue specificity:Constitutively expressed in cytotoxic T-cells.,
Background	fibrinogen like 2(FGL2) Homo sapiens The protein encoded by this gene is a secreted protein that is similar to the beta- and gamma-chains of fibrinogen. The carboxyl-terminus of the encoded protein consists of the fibrinogen-related domains (FRED). The encoded protein forms a tetrameric complex which is stabilized by interchain disulfide bonds. This protein may play a role in physiologic functions at mucosal sites. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using Fgl2 Monoclonal Antibody