

CD7 Monoclonal Antibody

| Catalog No | YP-mAb-13983 |
|---------------------------|---|
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB |
| Gene Name | CD7 |
| Protein Name | T-cell antigen CD7 |
| Immunogen | The antiserum was produced against synthesized peptide derived from the Internal region of human CD7. AA range:51-100 |
| Specificity | CD7 Monoclonal Antibody detects endogenous levels of CD7 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-1:2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | CD7; T-cell antigen CD7; GP40; T-cell leukemia antigen; T-cell surface antigen Leu-9; TP41; CD7 |
| Observed Band | 25kD |
| Cell Pathway | Membrane; Single-pass type I membrane protein. |
| Tissue Specificity | Liver, Muscle, Spleen, |
| Function | function:Not yet known.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain.,subunit:Interacts with SECTM1., |
| Background | This gene encodes a transmembrane protein which is a member of the immunoglobulin superfamily. This protein is found on thymocytes and mature T cells. It plays an essential role in T-cell interactions and also in T-cell/B-cell interaction during early lymphoid development. [provided by RefSeq, Jul 2008], |
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |










Products Images



Western Blot analysis of various cells using CD7 Monoclonal Antibody