



Ox40 Monoclonal Antibody

This receptor has been shown to activate NF-kapMAB through its interaction wi		
Reactivity Human;Rat;Mouse; Applications WB Gene Name TNFRSF4 Protein Name Tumor necrosis factor receptor superfamily member 4 Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human TNFRSF4. AA range:131-180 Specificity Ox40 Monoclonal Antibody detects endogenous levels of Ox40 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TNFRSF4; TXGP1L; Tumor necrosis factor receptor superfamily member 4; ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1 receptor; CD134 Observed Band 30kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain, Function function:Receptor for TNFSF4/OX40L/GP34, similarity:Contains 4 TNFR-Cys repeats., subunit:Interacts with TRAF2, TRAF3 and TRAF5. Background This receptor has been shown to activate NF-kapMAB through its interaction wit adaptor proteins TRAF2. Thockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2!L1.IBG.L2-XL, and thus suppresses apoptosis. The knockout studies in mice suggested that the receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul Cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul Cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul Cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul Cell-dependent B cell proliferation and differentiation.	Catalog No	YP-mAb-14064
Applications WB Gene Name TNFRSF4 Protein Name Tumor necrosis factor receptor superfamily member 4 Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human TNFRSF4. AA range:131-180 Specificity Ox40 Monoclonal Antibody detects endogenous levels of Ox40 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TNFRSF4; TXGP1L; Tumor necrosis factor receptor superfamily member 4; ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1 receptor; CD134 Observed Band 30kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain, Function function:Receptor for TNFSF4/OX40L/GP34_similarity.Contains 4 TNFR-Cys repeats.,subunit:Interacts with TRAF2, TRAF3 and TRAF5. Background This receptor shown to activate NF-kapMAB through its interaction with adaptor proteins TRAF2 and TRAF5. Knockout studies in mockout studies in proteins TRAF2 and TRAF5. Knockout studies in a nember of the TNF-receptor sugerfamily This receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2LINEDECL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in Tcell-dependent B cell proliferation and differentiation. [provided by RefSeq. Juli	Isotype	IgG
Gene Name TNFRSF4 Protein Name Tumor necrosis factor receptor superfamily member 4 Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human TNFRSF4. AA range:131-180 Specificity Ox40 Monoclonal Antibody detects endogenous levels of Ox40 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TNFRSF4; TXGP1L; Tumor necrosis factor receptor superfamily member 4; ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1 receptor; CD134 Observed Band 30kD Cell Pathway Membrane; Single-pass type I membrane protein. Function function:Receptor for TNFSF4/OX40L/GP34.,similarity.Contains 4 TNFR-Cys repeats., subunit:Interacts with TRAF2, TRAF3 and TRAF5., Background The protein encoded by this gene is a member of the TNF-receptor superfamily This receptor is been shown to activate NF-kapMAB through its interaction with adaptor proteins TRAF2	Reactivity	Human;Rat;Mouse;
Immunogen	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human TNFRSF4. AA range:131-180 Specificity Ox40 Monoclonal Antibody detects endogenous levels of Ox40 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TNFRSF4; TXGP1L; Tumor necrosis factor receptor superfamily member 4; ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1 receptor; CD134 Observed Band 30kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain, Function function:Receptor for TNFSF4/OX40L/GP34, similarity:Contains 4 TNFR-Cys repeats, subunit:Interacts with TRAF2, TRAF3 and TRAF5. Background The protein encoded by this gene is a member of the TNF-receptor superfamilithis receptor promotes the expression of apoptosis inhibitors BCL2 and adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and	Gene Name	TNFRSF4
Internal region of human TNFRSF4. AA range:131-180 Specificity Ox40 Monoclonal Antibody detects endogenous levels of Ox40 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TNFRSF4; TXGP1L; Tumor necrosis factor receptor superfamily member 4; ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1 receptor; CD134 Observed Band 30kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain, Function function:Receptor for TNFSF4/OX40L/GP34. similarity:Contains 4 TNFR-Cys repeats., subunit:Interacts with TRAF2, TRAF3 and TRAF5. Background The protein encoded by this gene is a member of the TNF-receptor superfamil This receptor has been shown to activate NF-kapMAB through its interaction with adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2L1/IBCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+1 cell response, as well as in cell-religentation and differentiation. [provided by RefSeq, Jul	Protein Name	Tumor necrosis factor receptor superfamily member 4
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TNFRSF4; TXGP1L; Tumor necrosis factor receptor superfamily member 4; ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1 receptor; CD134 Observed Band 30kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain, Function function:Receptor for TNFSF4/OX40L/GP34.,similarity:Contains 4 TNFR-Cys repeats.,subunit:Interacts with TRAF2, TRAF3 and TRAF5. Background This receptor has been shown to activate NF-kapMAB through its interaction wit adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2lL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+1 cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jule 1]	Immunogen	
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TNFRSF4; TXGP1L; Tumor necrosis factor receptor superfamily member 4; ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1 receptor; CD134 Observed Band 30kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain, Function function:Receptor for TNFSF4/OX40L/GP34.,similarity:Contains 4 TNFR-Cys repeats.,subunit:Interacts with TRAF2, TRAF3 and TRAF5., Background The protein encoded by this gene is a member of the TNF-receptor superfamil This receptor has been shown to activate NF-kapMAB through its interaction will adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2Int/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell profileration and differentiation. [provided by RefSeq, Jul	Specificity	Ox40 Monoclonal Antibody detects endogenous levels of Ox40 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TNFRSF4; TXGP1L; Tumor necrosis factor receptor superfamily member 4; ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1 receptor; CD134 Observed Band 30kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain, Function function: Receptor for TNFSF4/OX40L/GP34., similarity: Contains 4 TNFR-Cys repeats., subunit: Interacts with TRAF2, TRAF3 and TRAF5., The protein encoded by this gene is a member of the TNF-receptor superfamily This receptor has been shown to activate NF-kapMAB through its interaction will adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TNFRSF4; TXGP1L; Tumor necrosis factor receptor superfamily member 4; ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1 receptor; CD134 Observed Band 30kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain, Function function:Receptor for TNFSF4/OX40L/GP34., similarity:Contains 4 TNFR-Cys repeats., subunit:Interacts with TRAF2, TRAF3 and TRAF5. Background This receptor has been shown to activate NF-kapMAB through its interaction with adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2IL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jule 1] and the property of the provided by RefSeq, Jule 1] and the provided by RefSeq, Jule 2] and BCL2IL1 (PSCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jule 2] and BCL2IL1 (PSCL2-XL) and thus suppresses apoptosis.	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TNFRSF4; TXGP1L; Tumor necrosis factor receptor superfamily member 4; ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1 receptor; CD134 Observed Band 30kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain, Function function:Receptor for TNFSF4/OX40L/GP34.,similarity:Contains 4 TNFR-Cys repeats.,subunit:Interacts with TRAF2, TRAF3 and TRAF5., Background The protein encoded by this gene is a member of the TNF-receptor superfamil This receptor has been shown to activate NF-kapMAB through its interaction wit adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2IL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul	Purification	, , , , , , , , , , , , , , , , , , , ,
Purity ≥90% Storage Stability -20°C/1 year Synonyms TNFRSF4; TXGP1L; Tumor necrosis factor receptor superfamily member 4; ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1 receptor; CD134 Observed Band 30kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain, Function function:Receptor for TNFSF4/OX40L/GP34.,similarity:Contains 4 TNFR-Cys repeats.,subunit:Interacts with TRAF2, TRAF3 and TRAF5., Background This receptor has been shown to activate NF-kapMAB through its interaction will adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2lL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Juli	Dilution	WB 1:500-1:2000
Synonyms TNFRSF4; TXGP1L; Tumor necrosis factor receptor superfamily member 4; ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1 receptor; CD134 Observed Band Observed Band Membrane; Single-pass type I membrane protein. Fissue Specificity Brain, Function function:Receptor for TNFSF4/OX40L/GP34.,similarity:Contains 4 TNFR-Cys repeats.,subunit:Interacts with TRAF2, TRAF3 and TRAF5., The protein encoded by this gene is a member of the TNF-receptor superfamily This receptor has been shown to activate NF-kapMAB through its interaction wire adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2lL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jule 1]	Concentration	1 mg/ml
Synonyms TNFRSF4; TXGP1L; Tumor necrosis factor receptor superfamily member 4; ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1 receptor; CD134 Observed Band 30kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain, Function function:Receptor for TNFSF4/OX40L/GP34.,similarity:Contains 4 TNFR-Cys repeats.,subunit:Interacts with TRAF2, TRAF3 and TRAF5., Background The protein encoded by this gene is a member of the TNF-receptor superfamily This receptor has been shown to activate NF-kapMAB through its interaction will adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2IL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jule 1]	Purity	≥90%
ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1 receptor; CD134 Observed Band Observed Band Observed Band Observed Band Membrane; Single-pass type I membrane protein. Brain, Function function:Receptor for TNFSF4/OX40L/GP34.,similarity:Contains 4 TNFR-Cys repeats.,subunit:Interacts with TRAF2, TRAF3 and TRAF5., Background The protein encoded by this gene is a member of the TNF-receptor superfamily This receptor has been shown to activate NF-kapMAB through its interaction with adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2lL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul	Storage Stability	-20°C/1 year
Cell Pathway Membrane; Single-pass type I membrane protein. Function function:Receptor for TNFSF4/OX40L/GP34.,similarity:Contains 4 TNFR-Cys repeats.,subunit:Interacts with TRAF2, TRAF3 and TRAF5., Background The protein encoded by this gene is a member of the TNF-receptor superfamily This receptor has been shown to activate NF-kapMAB through its interaction with adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2lL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul	Synonyms	ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1
Tissue Specificity Brain, function: Receptor for TNFSF4/OX40L/GP34., similarity: Contains 4 TNFR-Cys repeats., subunit: Interacts with TRAF2, TRAF3 and TRAF5., The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor has been shown to activate NF-kapMAB through its interaction with adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2IL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul	Observed Band	30kD
Function function:Receptor for TNFSF4/OX40L/GP34.,similarity:Contains 4 TNFR-Cys repeats.,subunit:Interacts with TRAF2, TRAF3 and TRAF5., The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor has been shown to activate NF-kapMAB through its interaction with adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2IL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul	Cell Pathway	Membrane; Single-pass type I membrane protein.
repeats.,subunit:Interacts with TRAF2, TRAF3 and TRAF5., The protein encoded by this gene is a member of the TNF-receptor superfamily This receptor has been shown to activate NF-kapMAB through its interaction with adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2IL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul	Tissue Specificity	Brain,
This receptor has been shown to activate NF-kapMAB through its interaction wit adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that the receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2IL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul	Function	function:Receptor for TNFSF4/OX40L/GP34.,similarity:Contains 4 TNFR-Cys repeats.,subunit:Interacts with TRAF2, TRAF3 and TRAF5.,
	Background	BCL2IL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation. [provided by RefSeq, Jul



UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Ox40 Monoclonal Antibody