




CD72 Monoclonal Antibody

Catalog No	YP-mAb-14092
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	CD72
Protein Name	B-cell differentiation antigen CD72
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human CD72. AA range:170-220
Specificity	CD72 Monoclonal Antibody detects endogenous levels of CD72 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CD72; B-cell differentiation antigen CD72; Lyb-2; CD72
Observed Band	40kD
Cell Pathway	Membrane; Single-pass type II membrane protein.
Tissue Specificity	Pre-B-cells and B-cells but not terminally differentiated plasma cells.
Function	function:Plays a role in B-cell proliferation and differentiation. Associates with CD5.,online information:CD72,similarity:Contains 1 C-type lectin domain.,subunit:Homodimer; disulfide-linked.,tissue specificity:Pre-B-cells and B-cells but not terminally differentiated plasma cells.,
Background	function:Plays a role in B-cell proliferation and differentiation. Associates with CD5.,online information:CD72,similarity:Contains 1 C-type lectin domain.,subunit:Homodimer; disulfide-linked.,tissue specificity:Pre-B-cells and B-cells but not terminally differentiated plasma cells.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis of various cells using CD72 Monoclonal Antibody