

## CD296 Monoclonal Antibody

Catalog No         YP-mAb-14097           Isotype         IgG           Reactivity         Human;Mouse;Rat           Applications         WB           Gene Name         ART1           Protein Name         GPI-linked NAD(P)(+)—arginine ADP-ribosyltransferase 1           Immunogen         The antiserum was produced against synthesized peptide derived from the Internal region of human ART1. AN range:51-100           Specificity         CD296 Monoclonal Antibody detects endogenous levels of CD296 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dillution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         ART1; GPI-linked NAD(P)(+)—arginine ADP-ribosyltransferase 1; ADP-ribosyltransferase 22 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyltransferase 22 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyltransfer		
Reactivity         Human;Mouse;Rat           Applications         WB           Gene Name         ART1           Protein Name         GPI-linked NAD(P)(+)—arginine ADP-ribosyltransferase 1           Immunogen         The antiserum was produced against synthesized peptide derived from the Internal region of human ART1. AA range:51-100           Specificity         CD296 Monoclonal Antibody detects endogenous levels of CD296 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         ART1; GPI-linked NAD(P)(+)—arginine ADP-ribosyltransferase 1; ADP-ribosyltransferase C2 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl)-protein-Larginine = nicotinamide + N(ornega) (ADP-D-ribosyl)-protein-L-arginine = nicotinamide + N(ornega) (ADP-D-ribosyl)-protein-L-arginine, catalytic activity:NADP(+) + protein-L-arginine = nicotinamide + N(ornega) ((22-phospho-ADP)-D-ribosyl)-protein-L-arginine, similarity:Belongs to the A/rg-specific ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-piosylation is a posttranslational modification of proteins this interfered with by a variety of ba	Catalog No	YP-mAb-14097
Applications WB Gene Name ART1 Protein Name GPI-linked NAD(P)(+)arginine ADP-ribosyltransferase 1 Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human ART1. AA range:51-100 Specificity CD296 Monoclonal Antibody detects endogenous levels of CD296 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms ART1; GPI-linked NAD(P)(+)arginine ADP-ribosyltransferase 1; ADP-ribosyltransferase C2 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl))-protein-Larginine, actalytic activity: NADP(+) + N(omega) -(ADP-D-ribosyl)-protein-L-arginine = nicotinamide + N(omega) -(ADP-D-ribosyl)-protein-L-arginine, catalytic activity: NADP(+) + protein-L-arginine = nicotinamide + N(omega) -(ADP-D-ribosyl)-protein-L-arginine, catalytic activity: NADP(+) + protein-L-arginine, catalytic activity: NADP(+) + protein-L-arginine, catalytic activity: NADP(+) + N(omega) -(ADP-D-ribosyl)-protein-L-arginine = nicotinamide + N(omega) -(ADP-D-prosyl)-protein-L-arginine, catalytic activity: NADP(+) + protein-L-arginine = nicotinamide + N(omega) -(ADP-D-prosyl)-protein-L-arginine = nicotinamide + N(omega) - N(omega) - N(omega) - N(omega) - N(omega) - N(omega) - N(omega)	Isotype	IgG
Gene Name         ART1           Protein Name         GPI-linked NAD(P)(+)arginine ADP-ribosyltransferase 1           Immunogen         The antiserum was produced against synthesized peptide derived from the Internal region of human ART1. AA range:51-100           Specificity         CD296 Monoclonal Antibody detects endogenous levels of CD296 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         ART1; GPI-linked NAD(P)(+)arginine ADP-ribosyltransferase 1; ADP-ribosyltransferase C2 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl)transferase C2 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl)transferase 1; CD296           Observed Band         37kD           Cell Pathway         Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor.           Function         catalytic activity: NAD(+) + protein-L-arginine_, catalytic activity: NADP(+) + protein-L-arginine_, cataly	Reactivity	Human;Mouse;Rat
Protein Name         GPI-linked NAD(P)(+)arginine ADP-ribosyltransferase 1           Immunogen         The antiserum was produced against synthesized peptide derived from the Internal region of human ART1. AA range:51-100           Specificity         CD296 Monoclonal Antibody detects endogenous levels of CD296 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, lgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         ART1; GPI-linked NAD(P)(+)arginine ADP-ribosyltransferase 1; ADP-ribosyltransferase C2 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl)transferase 1; CD296           Observed Band         37kD           Cell Pathway         Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor.           Tissue Specificity         Skeletal muscle,           Function         catalytic activity:NAD(+) + protein-L-arginine_,catalytic activity:NADP(+) + protein-L-arginine_,icalalytic activity:NaDP(+) + protein-L-arginine_,ic	Applications	WB
Immunogen         The antiserum was produced against synthesized peptide derived from the Internal region of human ART1. AA range:51-100           Specificity         CD296 Monoclonal Antibody detects endogenous levels of CD296 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, lgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         ART1; GPI-linked NAD(P)(+)—arginine ADP-ribosyltransferase 1; ADP-ribosyltransferase 22 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl) transferase 1; CD296           Observed Band         37kD           Cell Pathway         Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor.           Tissue Specificity         Skeletal muscle,           Function         catalytic activity:NAD(+) + protein-L-arginine = nicotinamide + N(omega)-(ADP-D-ribosyl)-protein-L-arginine, catalytic activity:NADP(+) + protein-L-arginine = nicotinamide + N(omega)-(ADP-D-ribosyltransferase catalyzes the ADP-ribosyltain of arginine residues in proteins. Mono-ADP-ribosyltainsferase tamily.           Background         ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-r	Gene Name	ART1
Internal region of human ART1. AA range:51-100  Specificity  CD296 Monoclonal Antibody detects endogenous levels of CD296 protein.  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-1:2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  ART1; GPI-linked NAD(P)(+)—arginine ADP-ribosyltransferase 1; ADP-ribosyltransferase C2 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl)transferase 1; CD296  Observed Band  37kD  Cell Pathway  Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor.  Tissue Specificity  Skeletal muscle,  Function  catalytic activity:NAD(+) + protein-L-arginine = nicotinamide + N(omega)-(ADP-D-ribosyl)-protein-L-arginine.,catalytic activity:NADP(+) + protein-L-arginine.,catalytic activity:NADP(+) + protein-L-arginine.,catalytic activity:NADP(+) + protein-L-arginine = nicotinamide + N(omega)-((2-phospho-ADP)-D-ribosyl)-protein-L-arginine.,similarity:Belongs to the Arg-specific ADP-ribosyltransferase family.,  ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins this interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of E. coli. The amino acid sequence consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of glycosylphosphaltidylinositol (GPI)-anchored proteins. This gene was previously	Protein Name	GPI-linked NAD(P)(+)arginine ADP-ribosyltransferase 1
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms ART1; GPI-linked NAD(P)(+)-arginine ADP-ribosyltransferase 1; ADP-ribosyltransferase C2 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl)transferase 1; CD296  Observed Band 37kD  Cell Pathway Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor.  Tissue Specificity Skeletal muscle,  Function catalytic activity:NAD(+) + protein-L-arginine = nicotinamide + N(omega)-(ADP-D-ribosyl)-protein-L-arginine.,catalytic activity:NADP(+) + protein-L-arginine = nicotinamide + N(omega)-(2'-phospho-ADP)-D-ribosyly)-protein-L-arginine.,similarity:Belongs to the Arg-specific ADP-ribosyltransferase family.  Background ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosyltransferase family.  ADP-ribosyltransferase catalyzes the ADP-ribosylation of proteins this interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of £. coli. The amino acid sequence consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of glycosylposyphatioylinosity (GPI)-annotored proteins. This gene was previously	Immunogen	
Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         ART1; GPI-linked NAD(P)(+)arginine ADP-ribosyltransferase 1; ADP-ribosyltransferase C2 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl)transferase 1; CD296           Observed Band         37kD           Cell Pathway         Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor.           Tissue Specificity         Skeletal muscle,           Function         catalytic activity:NAD(+) + protein-L-arginine = nicotinamide + N(omega)-(ADP-D-ribosyl)-protein-L-arginine., catalytic activity:NADP(+) + protein-L-arginine = nicotinamide + N(omega)-((2"-phospho-ADP)-D-ribosyl)-protein-L-arginine., similarity:Belongs to the Arg-specific ADP-ribosyltransferase family.,           Background         ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins this interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of £. coli. The amino acid sequence consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of plycosylphosphalidylinositol (GPI)-anchored proteins. This gene was previously	Specificity	CD296 Monoclonal Antibody detects endogenous levels of CD296 protein.
Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-1:2000  Concentration  1 mg/ml  ≥90%  Storage Stability  -20°C/1 year  Synonyms  ART1; GPI-linked NAD(P)(+)arginine ADP-ribosyltransferase 1; ADP-ribosyltransferase C2 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl)transferase C1; CD296  Observed Band  37kD  Cell Pathway  Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor.  Tissue Specificity  Skeletal muscle,  Function  Catalytic activity: NAD(+) + protein-L-arginine = nicotinamide + N(omega)-(ADP-D-ribosyl)-protein-L-arginine., catalytic activity: NADP(+) + protein-L-arginine = nicotinamide + N(omega)-((2-phospho-ADP)-D-ribosyl)-protein-L-arginine., similarity: Belongs to the Arg-specific ADP-ribosyltransferase family.  Background  ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins the interfered with by a variety of bacterial toxins including cholera, perfussis, and heat-labile enterotoxins of E. coli. The amino acid sequence consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of predominantly hydrophobic N- and C-terminal regions, which is characteristic of predominantly hydrophobic N- and C-terminal regions, which is characteristic of plycosylphosphatidylinostic (GPI)-anchored proteins. This gene was previously	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-1:2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  ART1; GPI-linked NAD(P)(+)arginine ADP-ribosyltransferase 1; ADP-ribosyltransferase C2 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl)transferase 1; CD296  Observed Band  37kD  Cell Pathway  Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor.  Tissue Specificity  Skeletal muscle,  Function  catalytic activity:NAD(+) + protein-L-arginine = nicotinamide + N(omega)-(ADP-D-ribosyl)-protein-L-arginine.,catalytic activity:NADP(+) + protein-L-arginine = nicotinamide + N(omega)-(2'-phospho-ADP)-D-ribosyl)-protein-L-arginine.,similarity:Belongs to the Arg-specific ADP-ribosyltransferase family.,  Background  ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins the is interfered with by a variety of bacterial toxin is including cholera, pertussis, and heat-labile enterotoxins of E. coli. The amino acid sequence consists of predominantly hydrophobic. N- and C-terminal regions, which is characteristic of glycosylphosphatidylinositol (GPI)-anchored proteins. This gene was previously	Source	Monoclonal, Mouse,lgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       ART1; GPI-linked NAD(P)(+)arginine ADP-ribosyltransferase 1; ADP-ribosyltransferase C2 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl)transferase 1; CD296         Observed Band       37kD         Cell Pathway       Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor.         Tissue Specificity       Skeletal muscle,         Function       catalytic activity:NAD(+) + protein-L-arginine = nicotinamide + N(omega)-(ADP-D-ribosyl)-protein-L-arginine.,catalytic activity:NADP(+) + protein-L-arginine = nicotinamide + N(omega)-((2'-phospho-ADP)-D-ribosyl)-protein-L-arginine.,similarity:Belongs to the Arg-specific ADP-ribosyltransferase family.,         Background       ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins that is interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of E. coli. The amino acid sequence consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of glycosylphosphatidylinositol (GPI)-anchored proteins. This gene was previously	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms ART1; GPI-linked NAD(P)(+)arginine ADP-ribosyltransferase 1; ADP-ribosyltransferase C2 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl)transferase 1; CD296  Observed Band 37kD  Cell Pathway Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor.  Tissue Specificity Skeletal muscle,  Function catalytic activity:NAD(+) + protein-L-arginine = nicotinamide + N(omega)-(ADP-D-ribosyl)-protein-L-arginine., catalytic activity:NADP(+) + protein-L-arginine = nicotinamide + N(omega)-((2'-phospho-ADP)-D-ribosyl)-protein-L-arginine., similarity:Belongs to the Arg-specific ADP-ribosyltransferase family.  Background ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins that is interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of E. coli. The amino acid sequence consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of glycosylphosphatidylinositol (GPI)-anchored proteins. This gene was previously	Dilution	WB 1:500-1:2000
Storage Stability  -20°C/1 year  ART1; GPI-linked NAD(P)(+)arginine ADP-ribosyltransferase 1; ADP-ribosyltransferase C2 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl)transferase 1; CD296  Observed Band  37kD  Cell Pathway  Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor.  Tissue Specificity  Skeletal muscle,  catalytic activity:NAD(+) + protein-L-arginine = nicotinamide + N(omega)-(ADP-D-ribosyl)-protein-L-arginine.,catalytic activity:NADP(+) + protein-L-arginine = nicotinamide + N(omega)-((2'-phospho-ADP)-D-ribosyl)-protein-L-arginine.,similarity:Belongs to the Arg-specific ADP-ribosyltransferase family.,  ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins the is interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of E. coli. The amino acid sequence consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of glycosylphosphatidylinositol (GPI)-anchored proteins. This gene was previously	Concentration	1 mg/ml
ART1; GPI-linked NAD(P)(+)arginine ADP-ribosyltransferase 1; ADP-ribosyltransferase C2 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl)transferase 1; CD296  Observed Band 37kD  Cell Pathway Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor.  Tissue Specificity Skeletal muscle,  Function catalytic activity:NAD(+) + protein-L-arginine = nicotinamide + N(omega)-(ADP-D-ribosyl)-protein-L-arginine.,catalytic activity:NADP(+) + protein-L-arginine = nicotinamide + N(omega)-((2'-phospho-ADP)-D-ribosyl)-protein-L-arginine.,similarity:Belongs to the Arg-specific ADP-ribosyltransferase family.,  Background ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins that is interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of E. coli. The amino acid sequence consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of glycosylphosphatidylinositol (GPI)-anchored proteins. This gene was previously	Purity	≥90%
ADP-ribosyltransferase C2 and C3 toxin-like 1; ARTC1; Mono(ADP-ribosyl)transferase 1; CD296  Observed Band 37kD  Cell Pathway Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor.  Tissue Specificity Skeletal muscle,  Function catalytic activity:NAD(+) + protein-L-arginine = nicotinamide + N(omega)-(ADP-D-ribosyl)-protein-L-arginine.,catalytic activity:NADP(+) + protein-L-arginine = nicotinamide + N(omega)-((2'-phospho-ADP)-D-ribosyl)-protein-L-arginine.,similarity:Belongs to the Arg-specific ADP-ribosyltransferase family.,  Background ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins that is interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of E. coli. The amino acid sequence consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of glycosylphosphatidylinositol (GPI)-anchored proteins. This gene was previously	Storage Stability	-20°C/1 year
Cell Pathway  Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor.  Skeletal muscle,  Catalytic activity:NAD(+) + protein-L-arginine = nicotinamide + N(omega)-(ADP-D-ribosyl)-protein-L-arginine.,catalytic activity:NADP(+) + protein-L-arginine = nicotinamide + N(omega)-((2'-phospho-ADP)-D-ribosyl)-protein-L-arginine.,similarity:Belongs to the Arg-specific ADP-ribosyltransferase family.,  Background  ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins that is interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of E. coli. The amino acid sequence consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of glycosylphosphatidylinositol (GPI)-anchored proteins. This gene was previously	Synonyms	ADP-ribosyltransferase C2 and C3 toxin-like 1; ARTC1;
Tissue Specificity  Skeletal muscle,  catalytic activity:NAD(+) + protein-L-arginine = nicotinamide + N(omega)-(ADP-D-ribosyl)-protein-L-arginine.,catalytic activity:NADP(+) + protein-L-arginine = nicotinamide + N(omega)-((2'-phospho-ADP)-D-ribosyl)-protein-L-arginine.,similarity:Belongs to the Arg-specific ADP-ribosyltransferase family.,  ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins that is interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of E. coli. The amino acid sequence consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of glycosylphosphatidylinositol (GPI)-anchored proteins. This gene was previously	Observed Band	37kD
Function  catalytic activity:NAD(+) + protein-L-arginine = nicotinamide + N(omega)-(ADP-D-ribosyl)-protein-L-arginine.,catalytic activity:NADP(+) + protein-L-arginine = nicotinamide + N(omega)-((2'-phospho-ADP)-D-ribosyl)-protein-L-arginine.,similarity:Belongs to the Arg-specific ADP-ribosyltransferase family.,  ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins that is interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of E. coli. The amino acid sequence consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of glycosylphosphatidylinositol (GPI)-anchored proteins. This gene was previously	Cell Pathway	Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor.
protein-L-arginine = nicotinamide + N(omega)-((2'-phospho-ADP)-D-ribosyl)-protein-L-arginine.,similarity:Belongs to the Arg-specific ADP-ribosyltransferase family.,  ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins that is interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of E. coli. The amino acid sequence consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of glycosylphosphatidylinositol (GPI)-anchored proteins. This gene was previously	Tissue Specificity	Skeletal muscle,
proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins that is interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of E. coli. The amino acid sequence consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of glycosylphosphatidylinositol (GPI)-anchored proteins. This gene was previously	Function	protein-L-arginine = nicotinamide + N(omega)-((2'-phospho-ADP)-D-ribosyl)-protein-L-arginine.,similarity:Belongs to
	Background	proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins that is interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of E. coli. The amino acid sequence consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of glycosylphosphatidylinositol (GPI)-anchored proteins. This gene was previously



## UpingBio technology Co.,Ltd






matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using CD296 Monoclonal Antibody