



PKA I α reg Monoclonal Antibody

Catalog No	YP-mAb-14925
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	PRKAR1A
Protein Name	cAMP-dependent protein kinase type I-alpha regulatory subunit
Immunogen	The antiserum was produced against synthesized peptide derived from human KAP0. AA range:271-320
Specificity	PKA I α reg Monoclonal Antibody detects endogenous levels of PKA I α reg protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	PRKAR1A; PKR1; PRKAR1; TSE1; cAMP-dependent protein kinase type I-alpha regulatory subunit; Tissue-specific extinguisher 1; TSE1
Observed Band	43kD
Cell Pathway	Cell membrane .
Tissue Specificity	Four types of regulatory chains are found: I-alpha, I-beta, II-alpha, and II-beta. Their expression varies among tissues and is in some cases constitutive and in others inducible.
Function	disease:Defects in PRKAR1A are the cause of Carney complex type 1 (CNC1) [MIM:160980]. CNC is a multiple neoplasia syndrome characterized by spotty skin pigmentation, cardiac and other myxomas, endocrine tumors, and psammomatous melanotic schwannomas.,disease:Defects in PRKAR1A are the cause of intracardiac myxoma [MIM:255960]. Inheritance is autosomal recessive.,disease:Defects in PRKAR1A are the cause of primary pigmented nodular adrenocortical disease type 1 (PPNAD1) [MIM:610489]. Primary pigmented nodular adrenocortical disease is a rare bilateral adrenal defect causing ACTH-independent Cushing syndrome. Macroscopic appearance of the adrenals is characteristic with small pigmented micronodules observed in the cortex. PPNAD1 is most often diagnosed in patients with Carney complex, but it can also be observed in patients without other manifestations or familial



history.,PTM:The pseudop

Background

cAMP is a signaling molecule important for a variety of cellular functions. cAMP exerts its effects by activating the cAMP-dependent protein kinase, which transduces the signal through phosphorylation of different target proteins. The inactive kinase holoenzyme is a tetramer composed of two regulatory and two catalytic subunits. cAMP causes the dissociation of the inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits. Four different regulatory subunits and three catalytic subunits have been identified in humans. This gene encodes one of the regulatory subunits. This protein was found to be a tissue-specific extinguisher that down-regulates the expression of seven liver genes in hepatoma x fibroblast hybrids. Mutations in this gene cause Carney complex (CNC). This gene can fuse to the RET protooncog


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PKA I α reg Monoclonal Antibody