





## PP2A-B56- δ Monoclonal Antibody

| Catalog No         | YP-mAb-14950  |
|--------------------|---|
| Isotype            | IgG   |
| Reactivity         | Human;Mouse;Rat   |
| Applications       | WB  |
| Gene Name          | PPP2R5D   |
| Protein Name       | Serine/threonine-protein phosphatase 2A 56 kDa regulatory subunit delta isoform   |
| Immunogen          | The antiserum was produced against synthesized peptide derived from human PPP2R5D. AA range:544-593   |
| Specificity        | PP2A-B56- $\delta$ $$ Monoclonal Antibody detects endogenous levels of PP2A-B56- $\delta$ protein.  |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source             | Monoclonal, Mouse,IgG   |
| Purification       | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  |
| Dilution           | WB 1:500-1:2000   |
| Concentration      | 1 mg/ml   |
| Purity             | ≥90%  |
| Storage Stability  | -20°C/1 year  |
| Synonyms           | PPP2R5D; Serine/threonine-protein phosphatase 2A 56 kDa regulatory subunit delta isoform; PP2A B subunit isoform B'-delta; PP2A B subunit isoform B56-delta; PP2A B subunit isoform PR61-delta; PP2A B subunit isoform R5-delta   |
| Observed Band      | 70kD  |
| Cell Pathway       | Cytoplasm. Nucleus. Nuclear in interphase, nuclear during mitosis.  |
| Tissue Specificity | Isoform Delta-2 is widely expressed. Isoform Delta-1 is highly expressed in brain.  |
| Function           | function:The B regulatory subunit might modulate substrate selectivity and catalytic activity, and also might direct the localization of the catalytic enzyme to a particular subcellular compartment.,induction:By retinoic acid; in neuroblastoma cell lines.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the phosphatase 2A regulatory subunit B56 family.,subcellular location:Nuclear in interphase, nuclear during mitosis.,subunit:PP2A consists of a common heterodimeric core enzyme, composed of a 36 kDa catalytic subunit (subunit C) and a 65 kDa constant regulatory subunit (PR65 or subunit A), that associates with a variety of regulatory subunits. Proteins that associate with the core dimer include three families of regulatory subunits B (the R2/B/PR55/B55, R3/B"/PR72/PR130/PR59 and R5/B'/B56 families), the 48 kDa variable regulatory subunit, viral proteins, |



## UpingBio technology Co.,Ltd

📞 Tel: 400-999-8863 🔛 Emall: UpingBio@163.com



**Background** 

The product of this gene belongs to the phosphatase 2A regulatory subunit B family. Protein phosphatase 2A is one of the four major Ser/Thr phosphatases, and it is implicated in the negative control of cell growth and division. It consists of a common heteromeric core enzyme, which is composed of a catalytic subunit and a constant regulatory subunit, that associates with a variety of regulatory subunits. The B regulatory subunit might modulate substrate selectivity and catalytic activity. This gene encodes a delta isoform of the regulatory subunit B56 subfamily. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using PP2A-B56-

δ Monoclonal Antibody