



PP2C ϵ Monoclonal Antibody

Catalog No	YP-mAb-14952
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	PPM1L
Protein Name	Protein phosphatase 1L
Immunogen	The antiserum was produced against synthesized peptide derived from human PPM1L. AA range:119-168
Specificity	PP2C ϵ Monoclonal Antibody detects endogenous levels of PP2C ϵ protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	PPM1L; PP2CE; Protein phosphatase 1L; Protein phosphatase 1-like; Protein phosphatase 2C isoform epsilon; PP2C-epsilon
Observed Band	42kD
Cell Pathway	Membrane ; Single-pass type I membrane protein .
Tissue Specificity	Ubiquitous. Highly expressed in heart, placenta, lung, liver, kidney and pancreas.
Function	catalytic activity:A phosphoprotein + H(2)O = a protein + phosphate.,cofactor:Binds 2 magnesium or manganese ions per subunit.,function:Acts as a suppressor of the SAPK signaling pathways by associating with and dephosphorylating MAP3K7/TAK1 and MAP3K5, and by attenuating the association between MAP3K7/TAK1 and MAP2K4 or MAP2K6.,similarity:Belongs to the PP2C family.,similarity:Contains 1 PP2C-like domain.,subunit:Interacts with MAP3K7/TAK1 (By similarity). Interacts with MAP3K5.,tissue specificity:Ubiquitous. Highly expressed in heart, placenta, lung, liver, kidney and pancreas.,
Background	The protein encoded by this gene is a magnesium or manganese-requiring phosphatase that is involved in several signaling pathways. The encoded protein downregulates apoptosis signal-regulating kinase 1, a protein that initiates a signaling cascade that leads to apoptosis when cells are subjected to cytotoxic



stresses. This protein also is an endoplasmic reticulum transmembrane protein that helps regulate ceramide transport from the endoplasmic reticulum to the Golgi apparatus. Finally, this gene may be involved in adiposity since it is upregulated in adipose tissues. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2015],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PP2C ϵ Monoclonal Antibody