



EGFL4 Monoclonal Antibody

Catalog No YP-mAb-15896 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name MEGF8 Protein Name Multiple epidermal growth factor-like domains protein 8 Immunogen The antiserum was produced against synthesized peptide derived from human MEGF8. AA range:1243-1292 Specificity EGFL4 Monoclonal Antibody detects endogenous levels of EGFL4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like domains protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane ; Single-pass type I membrane protein 8; Epidermal growth factor-like domains, similarity: Contains 4 laminin EGF-like domains, similarity: Contains 5 EGF-like domains,		
Reactivity Human;Rat;Mouse; Applications WB Gene Name MEGF8 Protein Name Multiple epidermal growth factor-like domains protein 8 Immunogen The antiserum was produced against synthesized peptide derived from human MEGF8. AA range:1243-1292 Specificity EGFL4 Monoclonal Antibody detects endogenous levels of EGFL4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain, Pancreas, Plasma, Function similarity: Contains 12 Kelch repeats, similarity: Contains 2 CUB domains, similarity: Contains 4 laminin EGF-like domains. similarity: Contains 5 EGF-like domains, similarity: Contains 5 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Selects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012].	Catalog No	YP-mAb-15896
Applications Gene Name MEGF8 Protein Name Multiple epidermal growth factor-like domains protein 8 Immunogen The antiserum was produced against synthesized peptide derived from human MEGF8. AA range:1243-1292 Specificity EGFL4 Monoclonal Antibody detects endogenous levels of EGFL4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats, similarity:Contains 2 CUB domains, similarity:Contains 4 laminin EGF-like domains. similarity:Contains 5 EGF-like domains, similarity:Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012].	Isotype	IgG
Gene Name MEGF8 Protein Name Multiple epidermal growth factor-like domains protein 8 Immunogen The antiserum was produced against synthesized peptide derived from human MEGF8. AA range:1243-1292 Specificity EGFL4 Monoclonal Antibody detects endogenous levels of EGFL4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain, Pancreas, Plasma, similarity: Contains 12 Kelch repeats, similarity: Contains 2 CUB domains, similarity: Contains 4 larminin EGF-like domains, similarity: Contains 5 EGF-like domains, similarity: Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012].	Reactivity	Human;Rat;Mouse;
Immunogen	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human MEGF8. AA range:1243-1292 Specificity EGFL4 Monoclonal Antibody detects endogenous levels of EGFL4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain, Pancreas, Plasma, Function similarity: Contains 12 Kelch repeats, similarity: Contains 2 CUB domains, similarity: Contains 4 laminin EGF-like domains, similarity: Contains 5 EGF-like domains, similarity: Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a ca	Gene Name	MEGF8
Specificity EGFL4 Monoclonal Antibody detects endogenous levels of EGFL4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain, Pancreas, Plasma, Function similarity: Contains 12 Kelch repeats, similarity: Contains 2 CUB domains, similarity: Contains 4 laminin EGF-like domains, similarity: Contains 5 EGF-like domains, similarity: Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012],	Protein Name	Multiple epidermal growth factor-like domains protein 8
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane ; Single-pass type I membrane protein . Tissue Specificity Brain,Pancreas,Plasma, Function Similarity:Contains 12 Kelch repeats, similarity:Contains 2 CUB domains, similarity:Contains 5 EGF-like domains, similarity:Contains 7 PSI domains., similarity:Contains 5 EGF-like domains, similarity:Contains 7 PSI domains, Kelch repeats, and PSI domains. Defects in this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012], matters needing Avoid repeated freezing and thawing!	Immunogen	
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats, similarity:Contains 2 CUB domains, similarity:Contains 4 laminin EGF-like domains, similarity:Contains 5 EGF-like domains, similarity:Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012], matters needing Avoid repeated freezing and thawing!	Specificity	EGFL4 Monoclonal Antibody detects endogenous levels of EGFL4 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats, similarity:Contains 2 CUB domains, similarity:Contains 5 EGF-like domains, similarity:Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012], matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012], Matters needing Avoid repeated freezing and thawing!	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein . Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats, similarity:Contains 2 CUB domains, similarity:Contains 4 laminin EGF-like domains, similarity:Contains 5 EGF-like domains., similarity:Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012], matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats, similarity:Contains 2 CUB domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains., Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012], matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-1:2000
Storage Stability -20°C/1 year MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012], matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms MEGF8; C19orf49; EGFL4; KIAA0817; Multiple epidermal growth factor-like domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5 EGF-like domains., similarity:Contains 7 PSI domains., The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012], matters needing Avoid repeated freezing and thawing!	Purity	≥90%
domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth factor-like protein 4; EGF-like protein 4 Observed Band 255kD Cell Pathway Membrane; Single-pass type I membrane protein. Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains. Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012], matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Membrane ; Single-pass type I membrane protein . Tissue Specificity Brain,Pancreas,Plasma, Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 Iaminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains., Background The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012], matters needing Avoid repeated freezing and thawing!	Synonyms	domains protein 8; Multiple EGF-like domains protein 8; Epidermal growth
Tissue Specificity Brain,Pancreas,Plasma, Similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains., The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012], Matters needing Avoid repeated freezing and thawing!	Observed Band	255kD
Function similarity:Contains 12 Kelch repeats.,similarity:Contains 2 CUB domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains., The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012], matters needing Avoid repeated freezing and thawing!	Cell Pathway	Membrane ; Single-pass type I membrane protein .
domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5 EGF-like domains.,similarity:Contains 7 PSI domains., The protein encoded by this gene is a single-pass type I membrane protein of unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012], Matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Brain,Pancreas,Plasma,
unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2012], matters needing Avoid repeated freezing and thawing!	Function	domains.,similarity:Contains 4 laminin EGF-like domains.,similarity:Contains 5
	Background	unknown function that contains several EGF-like domains, Kelch repeats, and PSI domains. Defects in this gene are a cause of Carpenter syndrome 2. Two transcript variants encoding different isoforms have been found for this gene.
attention	matters needing attention	Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd






Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using EGFL4 Monoclonal Antibody