





IL-21 Monoclonal Antibody

Catalog No YP-mAb-15935 Isotype IgG Reactivity Human;Mouse Applications WB Gene Name IL21 Protein Name Interleukin-21 Immunogen The antiserum was produced against synthesized peptide derived from human iL21. AA range:91-140 Specificity IL-21 Monoclonal Antibody detects endogenous levels of IL-21 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms IL21; Interleukin-21; IL-21; Za11 Observed Band 18kD Cell Pathway Secreted Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function:Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1)		
Reactivity Human;Mouse Applications WB Gene Name IL21 Protein Name Interleukin-21 Immunogen The antiserum was produced against synthesized peptide derived from human IL21. AA range: 91-140 Specificity IL-21 Monoclonal Antibody detects endogenous levels of IL-21 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IL21; Interleukin-21; IL-21; Za11 Observed Band 18kD Cell Pathway Secreted . Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function:Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and adaptive immunity and play a role in proliferation of mature 3-band T-cells in response to activating simuli. In Supregy with IL15 and	Catalog No	YP-mAb-15935
Applications WB Gene Name IL21 Protein Name Interleukin-21 Immunogen The antiserum was produced against synthesized peptide derived from human IL21. AA range:91-140 Specificity IL-21 Monoclonal Antibody detects endogenous levels of IL-21 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse.lgG Puriffication The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms IL21; Interleukin-21; IL-21; Za11 Observed Band 18kD Cell Pathway Secreted. Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function:Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and sold in stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation. similarity Belongs to the IL-15/IL-21 family. lissue specificity, Expressed in activated of adaptive immunity. Indeed (1) (2) (1) in Survey with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation. similarity Belongs to the IL-15/IL-21 family. lissue specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells bu	Isotype	IgG
Gene Name IL21 Protein Name Interleukin-21 Immunogen The antiserum was produced against synthesized peptide derived from human IL21. AA range:91-140 Specificity IL-21 Monoclonal Antibody detects endogenous levels of IL-21 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IL21; Interleukin-21; IL-21; Za11 Observed Band 18kD Cell Pathway Secreted . Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function:Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL15. But yaregy with IL15 and NK cells and NK	Reactivity	Human;Mouse
Protein Name Interleukin-21 Immunogen The antiserum was produced against synthesized peptide derived from human IL21. AA range:91-140 Specificity IL-21 Monoclonal Antibody detects endogenous levels of IL-21 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IL21; Interleukin-21; IL-21; Za11 Observed Band 18kD Cell Pathway Secreted . Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function:Cytokine with immunoregulatory activity, May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and IN.15 and IL18 stimulates interferon gamma production in T-cells and IN.15 and IL18 stimulates interferon gamma production in T-cells and	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human IL21. AA range:91-140 Specificity IL-21 Monoclonal Antibody detects endogenous levels of IL-21 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IL21; Interleukin-21; IL-21; Za11 Observed Band 18kD Cell Pathway Secreted . Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function: Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (8y similarity). May play a role in proliferation of mature B- and T-cells (But in response to activating stimuli. In synergy with IL15, and IL18 as timulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation, similarity. Belongs to the IL-15fL-21 family, Lissue specificity: Expressed in activated CD4-positi	Gene Name	IL21
IL21. AA range:91-140 Specificity IL-21 Monoclonal Antibody detects endogenous levels of IL-21 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IL21; Interleukin-21; IL-21; Za11 Observed Band 18kD Cell Pathway Secreted . Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function:Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NkK) cells in synergy with L15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with I1.15 and I1.18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation, similarity:Belongs to the IL-15/IL-21 family, tissue specificity:Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Background Tries gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation.	Protein Name	Interleukin-21
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IL21; Interleukin-21; IL-21; Za11 Observed Band 18kD Cell Pathway Secreted. Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function: Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in poliferation and maturation of natural killer (NK) cells in synergy with IL15. May regulate proliferation of mature B- and T-cells in response to activating simuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation. similarity. Belongs to the IL-15/IL-21 family. Lissue specificity: Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Background This gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation,	Immunogen	
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IL21; Interleukin-21; IL-21; Za11 Observed Band 18kD Cell Pathway Secreted . Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function:Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL 15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation. similarity. Belongs to the IL-15/IL-21 family. tissue specificity: Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Background This gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentin	Specificity	IL-21 Monoclonal Antibody detects endogenous levels of IL-21 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml ≥90% Storage Stability -20°C/1 year Synonyms IL21; Interleukin-21; IL-21; Za11 Observed Band 18kD Cell Pathway Secreted . Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function:Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation, similarity. Belongs to the IL-15/IL-21 family, tissue specificity:Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. This gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immuner responses by inducing the differentiation.	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IL21; Interleukin-21; IL-21; Za11 Observed Band 18kD Cell Pathway Secreted . Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function: Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation.,similarity:Belongs to the IL-15/IL-21 family.,tissue specificity:Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. This gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation,	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IL21; Interleukin-21; IL-21; Za11 Observed Band 18kD Cell Pathway Secreted . Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function:Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of [gG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturationsimilarity.Belongs to the IL-15/IL-21 familytissue specificity:Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Background This gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation,	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms IL21; Interleukin-21; IL-21; Za11 Observed Band 18kD Cell Pathway Secreted . Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function: Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL 15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL 15 and IL 18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation., similarity:Belongs to the IL-15/IL-21 family., tissue specificity:Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Background This gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation,	Dilution	WB 1:500-1:2000
Storage Stability -20°C/1 year Synonyms IL21; Interleukin-21; IL-21; Za11 Observed Band 18kD Cell Pathway Secreted. Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function: Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation, similarity:Belongs to the IL-15/IL-21 family. tissue specificity:Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes., Background This gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation,	Concentration	1 mg/ml
Synonyms IL21; Interleukin-21; IL-21; Za11 Observed Band 18kD Cell Pathway Secreted . Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function:Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation.,similarity:Belongs to the IL-15/IL-21 family.,tissue specificity:Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes., Background This gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation,	Purity	≥90%
Cell Pathway Secreted . Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function:Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation., similarity:Belongs to the IL-15/IL-21 family., tissue specificity:Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes., Background This gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation,	Storage Stability	-20°C/1 year
Cell Pathway Secreted . Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function:Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation., similarity:Belongs to the IL-15/IL-21 family., tissue specificity:Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes., Background This gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation,	Synonyms	IL21; Interleukin-21; IL-21; Za11
Tissue Specificity Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes. Function function: Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation., similarity:Belongs to the IL-15/IL-21 family., tissue specificity:Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes., This gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation,	Observed Band	18kD
Function function:Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation., similarity:Belongs to the IL-15/IL-21 family., tissue specificity:Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes., This gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation,	Cell Pathway	Secreted .
between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation., similarity:Belongs to the IL-15/IL-21 family., tissue specificity:Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes., Background This gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation,	Tissue Specificity	
with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation,	Function	between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation., similarity:Belongs to the IL-15/IL-21 family., tissue specificity:Expressed in activated CD4-positive T-cells but not in CD8-positive
	Background	with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation,



UpingBio technology Co.,Ltd







killer cells, B cells and cytotoxic T cells. Dysregulation of this gene plays a role in multiple immune-mediated diseases including lupus, psoriasis and chronic inflammatory diseases. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Dec 2011],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using IL-21 Monoclonal Antibody