




CNG-2 Monoclonal Antibody

Catalog No	YP-mAb-16409
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CNGA2
Protein Name	Cyclic nucleotide-gated olfactory channel
Immunogen	The antiserum was produced against synthesized peptide derived from human CNGA2. AA range:391-440
Specificity	CNG-2 Monoclonal Antibody detects endogenous levels of CNG-2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CNGA2; CNCA; CNCA1; CNCG2; Cyclic nucleotide-gated olfactory channel; Cyclic nucleotide-gated cation channel 2; Cyclic nucleotide-gated channel alpha-2; CNG channel alpha-2; CNG-2; CNG2
Observed Band	83kD
Cell Pathway	Membrane; Multi-pass membrane protein.
Tissue Specificity	Testis,
Function	function:Odorant signal transduction is probably mediated by a G-protein coupled cascade using cAMP as second messenger. The olfactory channel can be shown to be activated by cyclic nucleotides which leads to a depolarization of olfactory sensory neurons.,similarity:Belongs to the cyclic nucleotide-gated cation channel (TC 1.A.1.5) family.,similarity:Contains 1 cyclic nucleotide-binding domain.,
Background	The protein encoded by this gene represents the alpha subunit of a cyclic nucleotide-gated olfactory channel. The encoded protein contains a carboxy-terminal leucine zipper that mediates channel formation. [provided by RefSeq, Jan 2010],
matters needing attention	Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using CNG-2 Monoclonal Antibody