






DNAM-1 Monoclonal Antibody

Catalog No	YP-mAb-17089
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	CD226
Protein Name	CD226 antigen
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human CD226. AA range:71-120
Specificity	DNAM-1 Monoclonal Antibody detects endogenous levels of DNAM-1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,lgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CD226; DNAM1; CD226 antigen; DNAX accessory molecule 1; DNAM-1; CD226
Observed Band	38kD
Cell Pathway	Cell membrane ; Single-pass type I membrane protein .
Tissue Specificity	Expressed by peripheral blood T-lymphocytes.
Function	function:Receptor involved in intercellular adhesion, lymphocyte signaling, cytotoxicity and lymphokine secretion mediated by cytotoxic T-lymphocyte (CTL) and NK cell.,PTM:Phosphorylated.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,subunit:Interacts with PVR and PVRL2.,tissue specificity:Expressed by peripheral blood T-lymphocytes.,
Background	This gene encodes a glycoprotein expressed on the surface of NK cells, platelets, monocytes and a subset of T cells. It is a member of the Ig-superfamily containing 2 Ig-like domains of the V-set. The protein mediates cellular adhesion of platelets and megakaryocytic cells to vascular endothelial cells. The protein also plays a role in megakaryocytic cell maturation. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2015],
matters needing attention	Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd








Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western Blot analysis of various cells using DNAM-1 Monoclonal Antibody