






TIFAB Mouse mAb


Catalog No	YP-mAb-18870
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	
Protein Name	
Immunogen	Synthetic peptide. This information is considered to be commercially sensitive.
Specificity	
Formulation	
Source	Monoclonal, Mouse, IgG
Purification	Affinity purification
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	18KDa
Calculated Molecular Weight	18KDa
Cell Pathway	
Tissue Specificity	
Function	
Background	TIFAB binds to TIFA (MIM 609028) and inhibits TIFA mediated NF kappa-B (NFKB1; MIM 164011) activation.
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western blot analysis of lysates from wild type (WT) and 293T cells transfected with TIFAB using [KD Validated] TIFAB Mouse mAb (A24024) at 1:2000 dilution incubated overnight at 4 °C. Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s



Western blot analysis of lysates from wild type (WT) and TIFAB knockdown (KD) MV-4-11 cells using [KD Validated] TIFAB Mouse mAb (A24024) at 1:2000 dilution incubated overnight at 4°C. Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.