





UQCRC2 mouse mAb

货号	YP-mAb-17993
同位型	lgG
应用	WB
种属	Human;Mouse;Rat
靶点	UQCRC2
基因名称	UQCRC2
蛋白名称	Cytochrome b-c1 complex subunit 2, mitochondrial (Complex III subunit 2) (Core protein II) (Ubiquinol-cytochrome-c reductase complex core protein 2)
免疫原	Synthesized peptide derived from human UQCRC2
特异性	This antibody detects endogenous levels of UQCRC2 at Human, Mouse,Rat
组成	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
来源	Monoclonal, mouse,lgG
稀释	WB 1:500-2000
纯化工艺	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
分子量	48kD
功能	Component of the ubiquinol-cytochrome c oxidoreductase, a multisubunit transmembrane complex that is part of the mitochondrial electron transport chain which drives oxidative phosphorylation. The respiratory chain contains 3 multisubunit complexes succinate dehydrogenase (complex II, CII), ubiquinol-cytochrome c oxidoreductase (cytochrome b-c1 complex, complex III, CIII) and cytochrome c oxidase (complex IV, CIV), that cooperate to transfer electrons derived from NADH and succinate to molecular oxygen, creating an electrochemical gradient over the inner membrane that drives transmembrane transport and the ATP synthase. The cytochrome b-c1 complex catalyzes electron transfer from ubiquinol to cytochrome c, linking this redox reaction to translocation of protons across the mitochondrial inner membrane, with protons being carried across the membrane as hydrogens on the quinol. In the proce
细胞定位	membrane protein ; Matrix side .
浓度	1 mg/ml
储存	-15°C to -25°C/1 year(Do not lower than -25°C)
有关注意事项	Avoid repeated freezing and thawing!
使用建议	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images

