







# MTDC mouse mAb

Catalog No	YP-mAb-11332
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	MTHFD2 NMDMC
Protein Name	MTDC
Immunogen	Synthesized peptide derived from human MTDC AA range: 121-171
Specificity	This antibody detects endogenous levels of MTDC at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, mouse,IgG
Purification	The antibody was affinity-purified from mouse serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Observed Band	
Calculated Molecular Weight	39kD
Cell Pathway	Mitochondrion.
Tissue Specificity	
Function	catalytic activity:5,10-methenyltetrahydrofolate + H(2)O = 10-formyltetrahydrofolate.,catalytic activity:5,10-methylenetetrahydrofolate + NAD(+) = 5,10-methenyltetrahydrofolate + NADH.,cofactor:Magnesium.,developmental stage:Expressed only in developing normal tissues.,miscellaneous:This NAD-dependent bifunctional enzyme has very different kinetic properties than the larger NADP-dependent trifunctional enzyme and is unique in that it requires formation of an enzyme-magnesium complex to allow binding of NAD.,similarity:Belongs to the tetrahydrofolate dehydrogenase/cyclohydrolase family.,subunit:Homodimer.,
Background	This gene encodes a nuclear-encoded mitochondrial bifunctional enzyme with methylenetetrahydrofolate dehydrogenase and methenyltetrahydrofolate cyclohydrolase activities. The enzyme functions as a homodimer and is unique in its absolute requirement for magnesium and inorganic phosphate. Formation of the enzyme-magnesium complex allows binding of NAD. Alternative splicing results in two different transcripts, one protein-coding and the other not



### UpingBio technology Co.,Ltd





protein-coding. This gene has a pseudogene on chromosome 7. [provided by RefSeq, Mar 2009],


## matters needing attention

Avoid repeated freezing and thawing!

#### **Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

### **Products Images**



Western Blot analysis of various cells using MTDC mouse mAb