




NAA40 Mouse mAb

| | |
|------------------------------------|---|
| Catalog No | YP-mAb-18896 |
| Isotype | IgG |
| Reactivity | Human,Mouse,Rat |
| Applications | WB |
| Gene Name | |
| Protein Name | |
| Immunogen | Recombinant protein (or fragment).This information is considered to be commercially sensitive. |
| Specificity | |
| Formulation | |
| Source | Monoclonal,Mouse,IgG |
| Purification | Affinity purification |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | NatD; NAT11; PATT1; hNatD; NAA40 |
| Observed Band | 27kDa |
| Calculated Molecular Weight | 27kDa |
| Cell Pathway | |
| Tissue Specificity | |
| Function | |
| Background | Enables H2A histone acetyltransferase activity; H4 histone acetyltransferase activity; and peptide-serine-N-acetyltransferase activity. Involved in N-terminal protein amino acid acetylation; histone H2A acetylation; and histone H4 acetylation. Located in centriolar satellite; cytosol; and nucleoplasm |
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |



Products Images



Western blot analysis of lysates from 293T cells, using NAA40 Mouse pAb (A7385) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.