



## XLF Rabbit pAb

|                           |   |
|---------------------------|---|
| <b>Catalog No</b>         | YP-Ab-18694   |
| <b>Isotype</b>            | IgG   |
| <b>Reactivity</b>         | Human, Mouse, Rat   |
| <b>Applications</b>       | WB  |
| <b>Gene Name</b>          | NHEJ1 XLF   |
| <b>Protein Name</b>       | Non-homologous end-joining factor 1 (Protein cernunnos) (XRCC4-like factor)   |
| <b>Immunogen</b>          | Synthesized peptide derived from human XLF  |
| <b>Specificity</b>        | This antibody detects endogenous levels of XLF at Human, Mouse,Rat  |
| <b>Formulation</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| <b>Source</b>             |   |
| <b>Purification</b>       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |
| <b>Dilution</b>           | WB 1:500-2000   |
| <b>Concentration</b>      | 1 mg/ml   |
| <b>Purity</b>             | ≥90%  |
| <b>Storage Stability</b>  | -20°C/1 year  |
| <b>Synonyms</b>           |   |
| <b>Observed Band</b>      | 33kD  |
| <b>Cell Pathway</b>       | Nucleus . Chromosome . Localizes to site of double-strand breaks; recruitment is dependent on XRCC5-XRCC6 (Ku) heterodimer. .   |
| <b>Tissue Specificity</b> | Ubiquitously expressed.   |
| <b>Function</b>           | DNA repair protein involved in DNA non-homologous end joining (NHEJ); required for double-strand break (DSB) repair and V(D)J recombination . Plays a key role in NHEJ by promoting the ligation of various mismatched and non-cohesive ends . Together with PAXX, collaborates with DNA polymerase lambda (POLL) to promote joining of non-cohesive DNA ends . May act in concert with XRCC5-XRCC6 (Ku) to stimulate XRCC4-mediated joining of blunt ends and several types of mismatched ends that are non-complementary or partially complementary . Associates with XRCC4 to form alternating helical filaments that bridge DNA and act like a bandage, holding together the broken DNA until it is repaired . The XRCC4-NHEJ1/XLF subcomplex binds to the DNA fragments of a DSB in a highly diffusive manner and robustly bridges two independent DNA molecules, holding the broken DNA fragments in close proximity to one other . The mobility of the bridges ensures that the ends remain accessible for further processing by other repair factors . Binds DNA in a length-dependent manner . |



## Background

### matters needing attention

Avoid repeated freezing and thawing!

### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images