

**(** Tel: 400-999-8863 **(** Email:3631691544@qq.com



## NADK Mouse mAb

Catalog No	YP-mAb-18591
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	
Protein Name	
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-180 of human NADK (NP_075394.3)
Specificity	
Formulation	
Source	
Purification	Affinity purification
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Cynonyms	dJ283E3.1; NADK
Observed Band	dJ283E3.1; NADK 49kDa
	·
Observed Band	·
Observed Band Cell Pathway	·
Observed Band Cell Pathway Tissue Specificity	A9kDa  NADK catalyzes the transfer of a phosphate group from ATP to NAD to generate
Observed Band Cell Pathway Tissue Specificity Function	49kDa
Observed Band Cell Pathway Tissue Specificity Function Background matters needing	A9kDa  NADK catalyzes the transfer of a phosphate group from ATP to NAD to generate NADP, which in its reduced form acts as an electron donor for biosynthetic reactions (Lerner et al., 2001 [PubMed 11594753])




## UpingBio technology Co.,Ltd

**(** Tel: 400-999-8863 **(** Email:3631691544@qq.com



## **Products Images**



Western blot analysis of various lysates using NADK Mouse pAb (A9111) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.