


YY1 Mouse mAb


货号	YP-mAb-18368
同位型	IgG
应用	WB
种属	Human, Mouse,Rat
其他名称	DELTA; NF-E1; UCRBP; GADEV5; INO80S; YIN-YANG-1; YY1
免疫原	A synthetic peptide corresponding to a sequence within amino acids 200-300 of human YY1 (P25490).
稀释	WB 1:1000 - 1:6000
纯化工艺	Affinity purification
分子量	60kDa
背景	YY1 is a ubiquitously distributed transcription factor belonging to the GLI-Kruppel class of zinc finger proteins. The protein is involved in repressing and activating a diverse number of promoters. YY1 may direct histone deacetylases and histone acetyltransferases to a promoter in order to activate or repress the promoter, thus implicating histone modification in the function of YY1.
浓度	1 mg/ml
储存	-15°C to -25°C/1 year(Do not lower than -25°C)
有关注意事项	Avoid repeated freezing and thawing!
使用建议	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunoprecipitation analysis of 300 µg extracts of HeLa cells using 3 µg YY1 antibody (A19569). Western

blot was performed from the immunoprecipitate using YY1 antibody (A19569) at a dilution of 1:2000



Western blot analysis of various lysates using YY1 Mouse mAb (A19569) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Western blot analysis of various lysates using YY1 Mouse mAb (A19569) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.