








## FOXO1 Mouse mAb


货号	YP-mAb-18370
同位型	IgG
应用	WB
种属	Human, Mouse,Rat
其他名称	FKH1; FKHR; FOXO1A
免疫原	Recombinant fusion protein containing a sequence corresponding to amino acids 450-650 of human FOXO1 (NP_002006.2).
稀释	WB 1:500 - 1:1000
纯化工艺	Affinity purification
分子量	70kDa
背景	This gene belongs to the forkhead family of transcription factors which are characterized by
	a distinct forkhead domain. The specific function of this gene has not yet been determined;
	however, it may play a role in myogenic growth and differentiation. Translocation of this
	gene with PAX3 has been associated with alveolar rhabdomyosarcoma
浓度	1 mg/ml
储存	Store at -20℃. Avoid freeze / thaw cycles.
	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
有关注意事项	Avoid repeated freezing and thawing!
使用建议	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**









Western blot analysis of lysates from HeLa cells using FOXO1 Mouse mAb (A25295) at 1:2000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Mouse


IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Western blot analysis of various lysates using FOXO1 Mouse mAb (A25295) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Mouse

IgG (H+L) (AS014) at 1:10000 dilution.

Lysates / proteins: 25 µg per lane.

Blocking buffer: 3 % nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 45s.