







RBM14 mouse mAb

Catalog No	YP-mAb-10076
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	RBM14 SIP
Protein Name	RNA-binding protein 14 (Paraspeckle protein 2) (PSP2) (RNA-binding motif protein 14) (RRM-containing coactivator activator/modulator) (Synaptotagmin-interacting protein) (SYT-interacting protein)
Immunogen	Synthesized peptide derived from human RBM14 AA range: 465-515
Specificity	This antibody detects endogenous levels of human RBM14
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, mouse,IgG
Purification	The antibody was affinity-purified from mouse serumby affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	74kD
Observed Band Cell Pathway	74kD Nucleus . Nucleus, nucleolus . Cytoplasm . In punctate subnuclear structures often located adjacent to splicing speckles, called paraspeckles (PubMed:11790299). Cytoplasmic localization is crucial for its function in suppressing the formation of aberrant centriolar protein complexes (PubMed:25385835).
	Nucleus . Nucleus, nucleolus . Cytoplasm . In punctate subnuclear structures often located adjacent to splicing speckles, called paraspeckles (PubMed:11790299). Cytoplasmic localization is crucial for its function in suppressing the formation of aberrant centriolar protein complexes
Cell Pathway	Nucleus . Nucleus, nucleolus . Cytoplasm . In punctate subnuclear structures often located adjacent to splicing speckles, called paraspeckles (PubMed:11790299). Cytoplasmic localization is crucial for its function in suppressing the formation of aberrant centriolar protein complexes (PubMed:25385835) Expressed in all tissues tested, including brain, heart, skeletal muscle, colon, thymus, spleen, kidney, liver, small intestine, placenta, lung and peripheral blood
Cell Pathway Tissue Specificity	Nucleus . Nucleus, nucleolus . Cytoplasm . In punctate subnuclear structures often located adjacent to splicing speckles, called paraspeckles (PubMed:11790299). Cytoplasmic localization is crucial for its function in suppressing the formation of aberrant centriolar protein complexes (PubMed:25385835) Expressed in all tissues tested, including brain, heart, skeletal muscle, colon, thymus, spleen, kidney, liver, small intestine, placenta, lung and peripheral blood
Cell Pathway Tissue Specificity Function	Nucleus . Nucleus, nucleolus . Cytoplasm . In punctate subnuclear structures often located adjacent to splicing speckles, called paraspeckles (PubMed:11790299). Cytoplasmic localization is crucial for its function in suppressing the formation of aberrant centriolar protein complexes (PubMed:25385835) Expressed in all tissues tested, including brain, heart, skeletal muscle, colon, thymus, spleen, kidney, liver, small intestine, placenta, lung and peripheral blood



UpingBio technology Co.,Ltd

€ Tel: 400-999-8863 🗷 Email:Upingbio.163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images