



GCET2 Mouse mAb

Catalog No	YP-mAb-18968
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	GCSAM GAL GCET2
Protein Name	Germinal center-associated signaling and motility protein (Germinal center B-cell-expressed transcript 2 protein) (Germinal center-associated lymphoma protein) (hGAL)
Immunogen	Synthesized peptide derived from human GCET2
Specificity	This antibody detects endogenous levels of GCET2 at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal,Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Calculated Molecular Weight	20kD
Cell Pathway	Cytoplasm. Cell membrane. It relocalizes from the cytoplasm to podosome-like structures upon cell treatment with IL6.
Tissue Specificity	Expressed in diffuse large B-cell lymphoma (DLBCL) and several germinal center (GC)-like lymphoma cell lines (at protein level). Highly expressed in normal GC lymphocytes and GC-derived malignancies. Expressed in thymus and spleen.
Function	Involved in the negative regulation of lymphocyte motility. It mediates the migration-inhibitory effects of IL6. Serves as a positive regulator of the RhoA signaling pathway. Enhancement of RhoA activation results in inhibition of lymphocyte and lymphoma cell motility by activation of its downstream effector ROCK. Is a regulator of B-cell receptor signaling, that acts through SYK kinase activation.
Background	This gene encodes a protein which may function in signal transduction pathways and whose expression is elevated in germinal cell lymphomas. It contains a putative PDZ-interacting domain, an immunoreceptor tyrosine-based activation motif (ITAM), and two putative SH2 binding sites. In B cells, its expression is



specifically induced by interleukin-4. Alternative splicing results in multiple transcript variants encoding different isoforms.

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images