








EFHD2 rabbit pAb

Catalog No	YP-Ab-11611
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	EFHD2 SWS1
Protein Name	EFHD2
Immunogen	Synthesized peptide derived from human EFHD2 AA range: 13-63
Specificity	This antibody detects endogenous levels of EFHD2 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	4 manufactural
Concentration	1 mg/ml
Purity	≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms Calculated	≥90% -20°C/1 year
Purity Storage Stability Synonyms Calculated Molecular Weight	≥90% -20°C/1 year 26kD
Purity Storage Stability Synonyms Calculated Molecular Weight Cell Pathway	≥90% -20°C/1 year 26kD Membrane raft . In a mouse immature B-cell line WEHI-231
Purity Storage Stability Synonyms Calculated Molecular Weight Cell Pathway Tissue Specificity	≥90% -20°C/1 year 26kD Membrane raft . In a mouse immature B-cell line WEHI-231 Found in lymphocytes; preferentially expressed in CD8+ cells. similarity:Contains 2 EF-hand domains.,tissue specificity:Found in lymphocytes;
Purity Storage Stability Synonyms Calculated Molecular Weight Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year 26kD Membrane raft . In a mouse immature B-cell line WEHI-231 Found in lymphocytes; preferentially expressed in CD8+ cells. similarity:Contains 2 EF-hand domains.,tissue specificity:Found in lymphocytes;
Purity Storage Stability Synonyms Calculated Molecular Weight Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year 26kD Membrane raft . In a mouse immature B-cell line WEHI-231 Found in lymphocytes; preferentially expressed in CD8+ cells. similarity:Contains 2 EF-hand domains.,tissue specificity:Found in lymphocytes; preferentially expressed in CD8+ cells.,



Products Images



Western blot analysis of lysates from HuvEc cells, primary antibody was diluted at 1:1000, 4° over night