





## ZNF22 Rabbit pAb

Catalog No	YP-Ab-18598
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	ZNF22 KOX15 KROX26
Protein Name	Zinc finger protein 22 (Zinc finger protein KOX15) (Zinc finger protein Krox-26)
Immunogen	Synthesized peptide derived from human ZNF22
Specificity	This antibody detects endogenous levels of ZNF22 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	25kD
0 11 0 41	
Cell Pathway	Nucleus .
Tissue Specificity	Nucleus .  In the embryo, expressed in developing craniofacial structures including dental epithelium of maxillary molar tooth organs, tongue epithelium and muscle, and craniofacial bone osteoblasts. In the adult, expressed in mesoderm-derived tissues such as skeletal muscle, heart, kidney and liver. Intermediate expression in spleen, thymus and brain. Low levels in endoderm-derived tissues such as intestine and colon.
-	In the embryo, expressed in developing craniofacial structures including dental epithelium of maxillary molar tooth organs, tongue epithelium and muscle, and craniofacial bone osteoblasts. In the adult, expressed in mesoderm-derived tissues such as skeletal muscle, heart, kidney and liver. Intermediate expression in spleen, thymus and brain. Low levels in endoderm-derived tissues such as intestine and colon.  Binds DNA through the consensus sequence 5'-CAATG-3'. May be involved in
Tissue Specificity	In the embryo, expressed in developing craniofacial structures including dental epithelium of maxillary molar tooth organs, tongue epithelium and muscle, and craniofacial bone osteoblasts. In the adult, expressed in mesoderm-derived tissues such as skeletal muscle, heart, kidney and liver. Intermediate expression in spleen, thymus and brain. Low levels in endoderm-derived tissues such as intestine and colon.
Tissue Specificity  Function	In the embryo, expressed in developing craniofacial structures including dental epithelium of maxillary molar tooth organs, tongue epithelium and muscle, and craniofacial bone osteoblasts. In the adult, expressed in mesoderm-derived tissues such as skeletal muscle, heart, kidney and liver. Intermediate expression in spleen, thymus and brain. Low levels in endoderm-derived tissues such as intestine and colon.  Binds DNA through the consensus sequence 5'-CAATG-3'. May be involved in
Tissue Specificity  Function  Background  matters needing	In the embryo, expressed in developing craniofacial structures including dental epithelium of maxillary molar tooth organs, tongue epithelium and muscle, and craniofacial bone osteoblasts. In the adult, expressed in mesoderm-derived tissues such as skeletal muscle, heart, kidney and liver. Intermediate expression in spleen, thymus and brain. Low levels in endoderm-derived tissues such as intestine and colon.  Binds DNA through the consensus sequence 5'-CAATG-3'. May be involved in transcriptional regulation and may play a role in tooth formation (By similarity).



## UpingBio technology Co.,Ltd

**C** Tel: 400-999-8863 **■** Email:3631691544@qq.com



Products Images