






ARIH1 Mouse mAb

Catalog No	YP-mAb-19040
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	
Protein Name	
Immunogen	Synthetic peptide. This information is considered to be commercially sensitive.
Specificity	
Formulation	
Source	Monoclonal,Mouse,IgG
Purification	Affinity purification
Dilution	WB 1:500-2000
Concentration	1 mg/ml
	. 224
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Storage Stability	-20°C/1 year
Storage Stability Synonyms Observed Band Calculated Molecular	-20°C/1 year ARI; HARI; HHARI; UBCH7BP; ARIH1
Storage Stability Synonyms Observed Band	-20°C/1 year ARI; HARI; HHARI; UBCH7BP; ARIH1 64kDa
Storage Stability Synonyms Observed Band Calculated Molecular Weight	-20°C/1 year ARI; HARI; HHARI; UBCH7BP; ARIH1 64kDa
Storage Stability Synonyms Observed Band Calculated Molecular Weight Cell Pathway	-20°C/1 year ARI; HARI; HHARI; UBCH7BP; ARIH1 64kDa
Storage Stability Synonyms Observed Band Calculated Molecular Weight Cell Pathway Tissue Specificity	-20°C/1 year ARI; HARI; HHARI; UBCH7BP; ARIH1 64kDa
Storage Stability Synonyms Observed Band Calculated Molecular Weight Cell Pathway Tissue Specificity Function	-20°C/1 year ARI; HARI; HHARI; UBCH7BP; ARIH1 64kDa 64kDa Enables enzyme binding activity; ubiquitin-protein transferase activity; and zinc ion binding activity. Involved in protein ubiquitination. Located in Lewy body; cytoplasm; and nuclear body. Colocalizes with cullin-RING ubiquitin ligase
Storage Stability Synonyms Observed Band Calculated Molecular Weight Cell Pathway Tissue Specificity Function Background matters needing	-20°C/1 year ARI; HARI; HHARI; UBCH7BP; ARIH1 64kDa 64kDa Enables enzyme binding activity; ubiquitin-protein transferase activity; and zinc ion binding activity. Involved in protein ubiquitination. Located in Lewy body; cytoplasm; and nuclear body. Colocalizes with cullin-RING ubiquitin ligase complex.



Products Images



Western blot analysis of various lysates using ARIH1 Mouse mAb (A17123) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: