


## LRRN3 rabbit pAb


| Catalog No   | YP-Ab-11292  |
|--|--|
| Isotype  | IgG  |
| Reactivity   | Human; Mouse;Rat   |
| Applications   | WB;IHC   |
| Gene Name  | LRRN3 Nbla10363 UNQ194/PRO220  |
| Protein Name   | LRRN3  |
| Immunogen  | Synthesized peptide derived from human LRRN3 AA range: 225-275   |
| Specificity  | This antibody detects endogenous levels of LRRN3 at Human/Mouse/Rat  |
| Formulation  | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source   | Polyclonal, Rabbit,IgG   |
| Purification   | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  |
| Dilution   | WB 1:500-2000;IHC-p 1:50-300   |
| Concentration  | 1 mg/ml  |
| Purity   | ≥90%   |
|  | 20076  |
| Storage Stability  | -20°C/1 year   |
| •  |  |
| Storage Stability  |  |
| Storage Stability Synonyms Calculated Molecular  | -20°C/1 year   |
| Storage Stability Synonyms Calculated Molecular Weight   | -20°C/1 year 78kD  |
| Storage Stability Synonyms Calculated Molecular Weight Cell Pathway  | -20°C/1 year 78kD  |
| Storage Stability Synonyms Calculated Molecular Weight Cell Pathway Tissue Specificity                                     | -20°C/1 year  78kD  Membrane ; Single-pass type I membrane protein .  similarity:Contains 1 fibronectin type-III domain.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 12 LRR (leucine-rich) |
| Storage Stability Synonyms Calculated Molecular Weight Cell Pathway Tissue Specificity Function                            | -20°C/1 year  78kD  Membrane ; Single-pass type I membrane protein .  similarity:Contains 1 fibronectin type-III domain.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 12 LRR (leucine-rich) |
| Storage Stability Synonyms Calculated Molecular Weight Cell Pathway Tissue Specificity Function Background matters needing | 78kD  Membrane; Single-pass type I membrane protein.  similarity:Contains 1 fibronectin type-III domain.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 12 LRR (leucine-rich) repeats.,       |



## **Products Images**



Western blot analysis of lysates from Jarkat cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for at 1900 retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).