







BPI Rabbit pAb

Catalog No	YP-Ab-19114
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	BPI
Protein Name	Bactericidal permeability-increasing protein (BPI) (CAP 57)
Immunogen	Synthesized peptide derived from human BPI
Specificity	This antibody detects endogenous levels of BPI at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
	2000/4
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20 C/T year
-	-20 C/T year
Synonyms	-20 C/T year 54kD
Synonyms Observed Band Calculated Molecular	
Synonyms Observed Band Calculated Molecular Weight	54kD Secreted . Cytoplasmic granule membrane . Membrane-associated in
Synonyms Observed Band Calculated Molecular Weight Cell Pathway	54kD Secreted . Cytoplasmic granule membrane . Membrane-associated in polymorphonuclear Leukocytes (PMN) granules
Synonyms Observed Band Calculated Molecular Weight Cell Pathway Tissue Specificity	Secreted . Cytoplasmic granule membrane . Membrane-associated in polymorphonuclear Leukocytes (PMN) granules Restricted to cells of the myeloid series. The cytotoxic action of BPI is limited to many species of Gram-negative bacteria; this specificity may be explained by a strong affinity of the very basic N-terminal half for the negatively charged lipopolysaccharides that are unique to the Gram-negative bacterial outer envelope. Has antibacterial activity against the Gram-negative bacterium P.aeruginosa, this activity is inhibited by LPS from
Synonyms Observed Band Calculated Molecular Weight Cell Pathway Tissue Specificity Function	Secreted . Cytoplasmic granule membrane . Membrane-associated in polymorphonuclear Leukocytes (PMN) granules Restricted to cells of the myeloid series. The cytotoxic action of BPI is limited to many species of Gram-negative bacteria; this specificity may be explained by a strong affinity of the very basic N-terminal half for the negatively charged lipopolysaccharides that are unique to the Gram-negative bacterial outer envelope. Has antibacterial activity against the Gram-negative bacterium P.aeruginosa, this activity is inhibited by LPS from



UpingBio technology Co.,Ltd







Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

