





## HSD17B7 Rabbit pAb

Catalog No	YP-Ab-19138
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	HSD17B7 UNQ2563/PRO6243
Protein Name	3-keto-steroid reductase (17-beta-hydroxysteroid dehydrogenase 7) (17-beta-HSD 7) (Estradiol 17-beta-dehydrogenase 7) (EC 1.1.1.62)
Immunogen	Synthesized peptide derived from human HSD17B7
Specificity	This antibody detects endogenous levels of HSD17B7 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Calculated Molecular Weight	38kD
Cell Pathway	Endoplasmic reticulum membrane ; Single-pass membrane protein .
Tissue Specificity	Highly expressed in adrenal gland, liver, lung and thymus. Expressed in breast, ovaries, pituitary gland, pregnant uterus, prostate, kidney, lymph node, small intestine, spinal cord and trachea. Weakly expressed in all other tissues tested.; [Isoform 3]: Expressed in eye ciliary epithelial cells and neuroendocrine cells.
Function	Bifunctional enzyme involved in steroid-hormone metabolism and cholesterol biosynthesis . Catalyzes the NADP(H)-dependent reduction of estrogens and androgens and regulates the biological potency of these steroids. Converts estrone (E1) to a more potent estrogen, 17beta-estradiol (E2) . Converts dihydrotestosterone (DHT) to its inactive form 5a-androstane-3b,17b-diol . Converts moderately progesterone to 3beta-hydroxypregn-4-ene-20-one, leading to its inactivation . Additionally, participates in the post-squalene cholesterol biosynthesis, as a 3-ketosteroid reductase . ; [Isoform 3]: Does not have enzymatic activities toward E1 and DHT.



## UpingBio technology Co.,Ltd







## **Background**

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

