



PDZK1 Rabbit pAb

Catalog No	YP-Ab-19149
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	PDZK1 CAP70 NHERF3 PDZD1
Protein Name	Na(+)/H(+) exchange regulatory cofactor NHE-RF3 (NHERF-3) (CFTR-associated protein of 70 kDa) (Na(+)/H(+) exchanger regulatory factor 3) (Na/Pi cotransporter C-terminal-associated protein 1) (NaPi-Cap1) (PDZ domain-containing protein 1) (Sodium-hydrogen exchanger regulatory factor 3)
Immunogen	Synthesized peptide derived from human PDZK1
Specificity	This antibody detects endogenous levels of SLC1A4 at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Calculated Molecular Weight	57kD
Cell Pathway	Membrane ; Peripheral membrane protein . Cell membrane . Associated with peripheral membranes. Localizes to the apical compartment of proximal tubular cells and to sinusoidal liver membranes. .
Tissue Specificity	Expression is limited to epithelial cells. Expressed in the kidney (brush border of proximal tubule), pancreas, liver, and small intestine. Expressed at a lower level in the adrenal cortex, testis and stomach. Overexpressed in breast, renal and lung carcinomas.
Function	A scaffold protein that connects plasma membrane proteins and regulatory components, regulating their surface expression in epithelial cells apical domains. May be involved in the coordination of a diverse range of regulatory processes for ion transport and second messenger cascades. In complex with SLC9A3R1, may cluster proteins that are functionally dependent in a mutual fashion and modulate the trafficking and the activity of the associated membrane proteins. May play a role in the cellular mechanisms associated with multidrug resistance through its



interaction with ABCC2 and PDZK1IP1. May potentiate the CFTR chloride channel activity. Required for normal cell-surface expression of SCARB1. Plays a role in maintaining normal plasma cholesterol levels via its effects on SCARB1. Plays a role in the normal localization and function of the chloride-anion exchanger SLC26A6 to the plasma membrane in the brush border of the proximal tubule of the kidney. May be involved in the regulation of proximal tubular Na(+)-dependent inorganic phosphate cotransport therefore playing an important role in tubule function (By similarity).

Background

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

