



UBC9 Monoclonal Antibody

Catalog No	YP-Ab-02341
Isotype	IgG
Reactivity	Human
Applications	WB;IHC;IF;FCM;ELISA
Gene Name	UBE2I
Protein Name	SUMO-conjugating enzyme UBC9
Immunogen	Purified recombinant fragment of human UBC9 expressed in E. Coli.
Specificity	UBC9 Monoclonal Antibody detects endogenous levels of UBC9 protein.
Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/200 - 1/1000. Immunofluorescence: 1/200 - 1/1000. Flow cytometry: 1/200 - 1/400. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	UBE2I; UBC9; UBCE9; SUMO-conjugating enzyme UBC9; SUMO-protein ligase; Ubiquitin carrier protein 9; Ubiquitin carrier protein I; Ubiquitin-conjugating enzyme E2 I; Ubiquitin-protein ligase I; p18
Observed Band	18kD
Cell Pathway	Nucleus . Cytoplasm . Cytoplasm, perinuclear region . Mainly nuclear (By similarity). In spermatocytes, localizes in synaptonemal complexes (PubMed:8610150). Recruited by BCL11A into the nuclear body (By similarity)..
Tissue Specificity	Expressed in heart, skeletal muscle, pancreas, kidney, liver, lung, placenta and brain. Also expressed in testis and thymus.
Function	catalytic activity:ATP + SUMO + protein lysine = AMP + diphosphate + protein N-SUMOyllysine..function:Accepts the ubiquitin-like proteins SUMO1, SUMO2, SUMO3 and SUMO4 from the UBLE1A-UBLE1B E1 complex and catalyzes their covalent attachment to other proteins with the help of an E3 ligase such as RANBP2 or CBX4. Essential for nuclear architecture and chromosome segregation..pathway:Protein modification; protein sumoylation..similarity:Belongs to the ubiquitin-conjugating enzyme family..subunit:Interacts with HIPK1, HIPK2 and PPM1J (By similarity). Forms a tight complex with RANGAP1 and RANBP2. Interacts with SIAH1 and PARP. Interacts with various transcription factors such as TCF3, TFAP2A, TFAP2B, TFAP2C, AR, ETS1 and SOX4. Interacts with human adenovirus E1A and human herpesvirus 6 IE2. Interacts with RWDD3; the interaction enhances the



sumoylation of a number of proteins such as HIF1A

Background


The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. Four alternatively spliced transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2008],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images


Western Blot analysis using UBC9 Monoclonal Antibody against HeLa (1), HepG2 (2), and Cos7 (3) cell lysate.



Immunohistochemistry analysis of paraffin-embedded prostate tissues with DAB staining using UBC9 Monoclonal Antibody.



Immunofluorescence analysis of HepG2 cells using UBC9 Monoclonal Antibody (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.



Flow cytometric analysis of HepG2 cells using UBC9 Monoclonal Antibody (green) and negative control (red).