



SNX6 Rabbit pAb

Catalog No	YP-Ab-19192
Isotype	IgG
Reactivity	Human,Mouse
Applications	WB
Gene Name	SNX6
Protein Name	Sorting nexin-6 (TRAF4-associated factor 2)
Immunogen	Synthesized peptide derived from human SNX6
Specificity	This antibody detects endogenous levels of SNX6 at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Calculated Molecular Weight	45kD
Cell Pathway	Early endosome . Early endosome membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasmic vesicle . Cytoplasm . Nucleus . Interaction with SNX1 or SNX2 promotes location at endosome membranes (PubMed:19935774). Only a minor proportion is seen in the nucleus. .
Tissue Specificity	
Function	Involved in several stages of intracellular trafficking. Interacts with membranes phosphatidylinositol 3,4-bisphosphate and/or phosphatidylinositol 4,5-bisphosphate (Probable). Acts in part as component of the retromer membrane-deforming SNX-BAR subcomplex . The SNX-BAR retromer mediates retrograde transport of cargo proteins from endosomes to the trans-Golgi network (TGN) and is involved in endosome-to-plasma membrane transport for cargo protein recycling. The SNX-BAR subcomplex functions to deform the donor membrane into a tubular profile called endosome-to-TGN transport carrier (ETC) (Probable). Does not have in vitro vesicle-to-membrane remodeling activity . Involved in retrograde endosome-to-TGN transport of lysosomal enzyme receptor IGF2R . May function as link between transport vesicles and dynactin (Probable).



Negatively regulates retrograde transport of BACE1 from the cell surface to the trans-Golgi network . Involved in E-cadherin sorting and degradation; inhibits PIP5K1C isoform 3-mediated E-cadherin degradation . In association with GIT1 involved in EGFR degradation. Promotes lysosomal degradation of CDKN1B (By similarity). May contribute to transcription regulation (Probable).

Background**matters needing attention**

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images