



# DHX15 Mouse mAb

<b>Catalog No</b>	YP-mAb-19198
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	DHX15 DBP1 DDX15
<b>Protein Name</b>	Putative pre-mRNA-splicing factor ATP-dependent RNA helicase DHX15 (ATP-dependent RNA helicase #46) (DEAH box protein 15)
<b>Immunogen</b>	Synthesized peptide derived from human DHX15
<b>Specificity</b>	This antibody detects endogenous levels of DHX15 at Human, Mouse
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal,Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Calculated Molecular Weight</b>	87kD
<b>Cell Pathway</b>	Nucleus . Nucleus, nucleolus .
<b>Tissue Specificity</b>	Ubiquitous.
<b>Function</b>	RNA helicase involved in mRNA processing and antiviral innate immunity . Pre-mRNA processing factor involved in disassembly of spliceosomes after the release of mature mRNA . In cooperation with TFIP11 seem to be involved in the transition of the U2, U5 and U6 snRNP-containing IL complex to the snRNP-free IS complex leading to efficient debranching and turnover of excised introns . Plays a key role in antiviral innate immunity by promoting both MAVS-dependent signaling and NLRP6 inflammasome . Acts as an RNA virus sensor: recognizes and binds viral double stranded RNA (dsRNA) and activates the MAVS-dependent signaling to produce interferon-beta and interferon lambda-3 (IFNL3) . Involved in intestinal antiviral innate immunity together with NLRP6: recognizes and binds viral dsRNA and promotes activation of the NLRP6 inflammasome in intestinal epithelial cells to restrict infection by enteric viruses . The NLRP6 inflammasome acts by promoting maturation and secretion of IL18 in



the extracellular milieu . Also involved in antibacterial innate immunity by promoting Wnt-induced antimicrobial protein expression in Paneth cells (By similarity).

## Background

### matters needing attention

Avoid repeated freezing and thawing!

### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images