



# CXXC1 Mouse mAb

<b>Catalog No</b>	YP-mAb-19246
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human,Mouse,Rat
<b>Applications</b>	WB
<b>Gene Name</b>	CXXC1 CFP1 CGBP PCCX1 PHF18
<b>Protein Name</b>	CpG-binding protein (CXXC-type zinc finger protein 1) (PHD finger and CXXC domain-containing protein 1)
<b>Immunogen</b>	Synthesized peptide derived from human CXXC1
<b>Specificity</b>	This antibody detects endogenous levels of CXXC1 at Human, Mouse
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal,Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Calculated Molecular Weight</b>	72kD
<b>Cell Pathway</b>	Nucleus speckle . Nucleus . Associated with euchromatin. During mitosis, excluded from condensed chromosomes.
<b>Tissue Specificity</b>	Ubiquitous.
<b>Function</b>	Transcriptional activator that exhibits a unique DNA binding specificity for CpG unmethylated motifs with a preference for CpGG.
<b>Background</b>	
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images