



NGAL mouse mAb

| | |
|------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Catalog No | YP-mAb-12478 |
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB |
| Gene Name | NGAL |
| Protein Name | NGAL |
| Immunogen | Synthesized peptide derived from human NGAL |
| Specificity | This antibody detects endogenous levels of Human NGAL |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-1:2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/ 1 year |
| Synonyms | Neutrophil gelatinase-associated lipocalin (NGAL;25 kDa alpha-2-microglobulin-related subunit of MMP-9;Lipocalin-2;Oncogene 24p3;Siderocalin LCN2;p25) |
| Calculated Molecular Weight | 22kD |
| Cell Pathway | Secreted . Cytoplasmic granule lumen . Cytoplasmic vesicle lumen . Upon binding to the SLC22A17 (24p3R) receptor, it is internalized (By similarity). Releases the bound iron in the acidic lumen of cytoplasmic vesicles (PubMed:12453413, PubMed:20581821). . |
| Tissue Specificity | Detected in neutrophils (at protein level) (PubMed:7683678, PubMed:8298140). Expressed in bone marrow and in tissues that are prone to exposure to microorganism. High expression is found in bone marrow as well as in uterus, prostate, salivary gland, stomach, appendix, colon, trachea and lung. Not found in the small intestine or peripheral blood leukocytes. |
| Background | |
| matters needing attention | Avoid repeated freezing and thawing! |

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images