# FAF2 rabbit pAb

| | |
|---|---|
| **Catalog No** | YP-Ab-09055 |
| **Isotype** | IgG |
| **Reactivity** | Human; Mouse;Rat |
| **Applications** | WB |
| **Gene Name** | FAF2 ETEA KIAA0887 UBXD8 UBXN3B |
| **Protein Name** | FAF2 |
| **Immunogen** | Synthesized peptide derived from human FAF2 AA range: 120-170 |
| **Specificity** | This antibody detects endogenous levels of FAF2 at Human/Mouse/Rat |
| **Formulation** | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| **Source** | Polyclonal, Rabbit,IgG |
| **Purification** | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| **Dilution** | WB 1：500-2000 |
| **Concentration** | 1 mg/ml |
| **Purity** | ≥90% |
| **Storage Stability** | -20°C/1 year |
| **Synonyms** | |
| **Calculated Molecular Weight** | 49kD |
| **Cell Pathway** | Cytoplasm . Lipid droplet . Endoplasmic reticulum . |
| **Tissue Specificity** | Broadly expressed, with highest levels in brain. |
| **Function** | caution:It seems odd to name this protein FAS-associated factor, but not have any mention of this in the CC lines.,miscellaneous:Up-regulated in T-cells and eosinophils from patients with atopic dermatitis.,similarity:Contains 1 UBA domain.,similarity:Contains 1 UBX domain.,tissue specificity:Broadly expressed, with highest levels in brain., |
| **Background** | The protein encoded by this gene is highly expressed in peripheral blood of patients with atopic dermatitis (AD), compared to normal individuals. It may play a role in regulating the resistance to apoptosis that is observed in T cells and eosinophils of AD patients. [provided by RefSeq, Jul 2008], |
| **matters needing attention** | Avoid repeated freezing and thawing! |

**Usage suggestions** This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images

Western blot analysis of lysates from HCT116 cells, primary antibody was diluted at 1:1000, 4°over night