



IL-17A Rabbit mAb

Catalog No	YP-rAb-16986
Isotype	IgG
Reactivity	Mouse
Applications	WB,IF,IP,ELISA
Gene Name	IL17A CTLA8 IL17
Protein Name	IL-17
Purification Process	Protein A
Specificity	Endogenous
Formulation	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source	Monoclonal, Rabbit,IgG
Dilution	WB 1:500-1:2000; IF 1:200-1:1000; ELISA 1:5000-1:20000; IP 1:50-1:200;
Concentration	0.5 mg/ml
Purity	≥90%
Storage Stability	-15° C to -25° C/1 year(Do not lower than -25° C)
Synonyms	Interleukin-17A ; IL-17 ; IL-17A ; Cytotoxic T-lymphocyte-associated antigen 8 ; CTLA-8 ;
Observed Band	18kD
Calculated Molecular Weight	18kD
Cell Pathway	Secreted .
Tissue Specificity	Expressed in memory Th17 cells (at protein level).
Function	regulation of transcription, DNA-dependent, regulation of transcription from RNA polymerase II promoter, protein amino acid glycosylation, apoptosis, defense response, inflammatory response, immune response, cell-cell signaling, cell death, glycoprotein metabolic process, glycoprotein biosynthetic process, response to wounding, positive regulation of biosynthetic process, positive regulation of macromolecule biosynthetic process, positive regulation of macromolecule metabolic process, positive regulation of gene expression, programmed cell death, death, positive regulation of cellular biosynthetic process, biopolymer glycosylation, regulation of transcription, positive regulation of transcription, DNA-dependent, positive regulation of nucleobase, nucleoside, nucleotide and nucleic acid metabolic process, positive regulation of transcription, positive regulation of transcription from RNA polymerase II promoter, positive regulation of nitrogen compound metabolic process, regulation of RNA metabolic

杭州臻优品生物科技有限公司

热销产品:

蛋白、一抗、抗体对、ELISA试剂盒、生化试剂盒
CCK8试剂盒、QPCR检测试剂盒

检测服务:

ELISA检测及定制服务 | 生化检测 | PCR、QPCR检测 | WB检测
| CO-IP检测 | 切片 | 染色 | 免疫组化 | 免疫荧光 | 透射电镜全套
| 宏基因组、转录组、基因组、蛋白组、代谢组测序



关注官网



关注客服



process, positive regulation of RNA metabolic process, glycosylation,

Background


function:Induces stromal cells to produce proinflammatory and hematopoietic cytokines. Enhances the surface expression of the intracellular adhesion molecule-1 (ICAM-1) in fibroblasts.,online information:Interleukin-17 entry,PTM:Found both in glycosylated and nonglycosylated forms.,similarity:Belongs to the IL-17 family.,subunit:Homodimer.,tissue specificity:Restricted to activated memory T-cells.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-IL-17A antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: RAW264.7 Lane 2: RAW264.7 was treated with LPS (0.1 μ g/ml) for 4 hou

