



FMR1 Rabbit mAb

Catalog No	YP-rAb-17212
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB,IHC,IF,ELISA
Gene Name	FMR1
Protein Name	Fragile X mental retardation 1 protein
Purification Process	Protein A
Specificity	Endogenous
Formulation	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source	Monoclonal, Rabbit,IgG
Dilution	IHC 1:200-1:1000; WB 1:2000-1:10000; IF 1:200-1:1000; ELISA 1:5000-1:20000; Note: For IHC, we suggest antigen retrieval with TE buffer pH 9.0
Concentration	0.5 mg/ml
Purity	≥90%
Storage Stability	-15° C to -25° C/1 year(Do not lower than -25° C)
Synonyms	FMR1 ; Fragile X mental retardation protein 1 ; FMRP ; Protein FMR-1
Observed Band	70-77kD
Calculated Molecular Weight	71kD
Cell Pathway	Nucleus . Nucleus, nucleolus . Chromosome, centromere . Chromosome . Cytoplasm . Cytoplasm, perinuclear region . Cytoplasm, Cytoplasmic ribonucleoprotein granule . Perikaryon . Cell projection, neuron projection . Cell projection, axon . Cell projection, dendrite . Cell projection, dendritic spine . Cell junction, synapse, synaptosome . Cell projection, growth cone . Cell projection, filopodium tip . Cell junction, synapse . Cell junction, synapse, postsynaptic cell membrane . Cell junction, synapse, presynaptic cell membrane . Cell membrane . Cytoplasm, Stress granule . Colocalizes with H2AX/H2A.x in pericentromeric heterochromatin in response to DNA damaging agents (By similarity). Localizes on meiotic pachytene-stage chromosomes (By similarity). Forms nuclear foci representing sites of ongoing DNA replication in response to DNA damaging agents (By similarity). Shuttles between nucleus and cytoplasm in a XPO1/CRM1-dependent manner (PubMed:10196376). Localizes to cytoplasmic ribonucleoprotein granules, also referred to as messenger ribonucleoprotein particles or mRNPs, along dendrites and dendritic spines (PubMed:9659908, PubMed:14532325). FMR1-containing cytoplasmic granules colocalize to F-actin-rich structures, including filopodium, spines and growth cone during the development of hippocampal neurons (By similarity). FMR1-containing

杭州臻优品生物科技有限公司

热销产品:

蛋白、一抗、抗体对、ELISA试剂盒、生化试剂盒
CCK8试剂盒、QPCR检测试剂盒

检测服务:

ELISA检测及定制服务 | 生化检测 | PCR、QPCR检测 | WB检测
ICO-IP检测 | 切片 | 染色 | 免疫组化 | 免疫荧光 | 透射电镜全套
| 宏基因组、转录组、基因组、蛋白组、代谢组测序



关注官网



关注客服



cytoplasmic granules are transported out of the soma along axon and dendrite to synaptic contacts in a microtubule- and kinesin-dependent manner (PubMed:12417734, PubMed:15380484). Colocalizes with CACNA1B in the cytoplasm and at the cell membrane of neurons (By similarity). Colocalizes with CYFIP1, CYFIP2, NXF2 and ribosomes in the perinuclear region (By similarity). Colocalizes with CYFIP1 and EIF4E in dendrites and probably at synapses (By similarity). Colocalizes with FXR1, kinesin, 60S acidic ribosomal protein RPLP0 and SMN in cytoplasmic granules in the soma and neurite cell processes (PubMed:12417734, PubMed:18093976, PubMed:16636078). Colocalizes with FXR1 and FXR2 in discrete granules, called fragile X granules (FXGs), along axon and presynaptic compartments (By similarity). Colocalizes with TDRD3 in cytoplasmic stress granules (SGs) in response to various cellular stress (PubMed:18632687, PubMed:18664458, PubMed:16636078). .; [Isoform 6]: Cytoplasm . Cytoplasm, perinuclear region .; [Isoform 9]: Cytoplasm .; [Isoform 10]: Nucleus . Nucleus, Cajal body . Colocalizes with Colin and SMN in Cajal bodies (PubMed:24204304).; [Isoform 11]: Nucleus . Nucleus, Cajal body .

Tissue Specificity

Expressed in the brain, cerebellum and testis (PubMed:8401578). Also expressed in epithelial tissues (PubMed:8401578). Expressed in mature oligodendrocytes (OLGs) (PubMed:23891804). Expressed in fibroblast (PubMed:24204304). Expressed in neurons, Purkinje cells and spermatogonias (at protein level) (PubMed:8401578). Expressed in brain, testis and placenta (PubMed:8504300). Expressed in neurons and lymphocytes (PubMed:8504300).

Function

Alternative products:At least 12 different isoforms are produced,Disease:Defects in FMR1 are the cause of fragile X syndrome. [MIM:300624]. It is a common genetic disease (has a prevalence of one in every 2000 children) which is characterized by moderate to severe mental retardation, macroorchidism (enlargement of the testicles), large ears, prominent jaw, and high-pitched, jocular speech. The defect in most fragile X syndrome patients results from an amplification of a CGG repeat region which is directly in front of the coding region.,Disease:Defects in FMR1 are the cause of fragile X tremor/ataxia syndrome (FXTAS) [MIM:300623]. In FXTAS, the expanded repeats range in size from 55 to 200 repeats and are referred to as 'premutations'. Full repeat expansions with greater than 200 repeats results in fragile X mental retardation syndrome [MIM:300624]. Carriers of the premutation typically do not show the full fragile X syndrome phenotype, but comprise a subgroup that may have some physical features of fragile X syndrome or mild cognitive and emotional problems.,Function:RNA-binding protein that plays a role in intracellular RNA transport and in the regulation of translation of target mRNAs. Associated with polysomes. May play a role in the transport of mRNA from the nucleus to the cytoplasm. Binds strongly to poly(G), binds moderately to poly(U) but shows very little binding to poly(A) or poly(C).,miscellaneous:RNA-binding activity is inhibited by RANBP9.,miscellaneous:The mechanism of the severe phenotype in the Asn-304 patient lies in the sequestration of bound mRNAs in nontranslatable mRNP particles. In the absence of FMRP, these same mRNAs may be partially translated via alternate mRNPs, although perhaps abnormally localized or regulated, resulting in typical fragile X syndrome. Asn-304 mutation maps to a position within the second KH domain of FMRP that is critical for stabilizing sequence-specific RNA-protein interactions. Asn-304 mutation abrogates the association of the FMRP KH 2 domain with its target, kissing complex RNA.,PTM:Phosphorylated on several serine residues.,similarity:Belongs to the FMR1 family.,similarity:Contains 1 KH domain.,similarity:Contains 2 KH domains.,subunit:Homooligomer. Found in a RNP granule complex with IGF2BP1. Directly interacts with SMN and TDRD3. Interacts with the SMN core complex that contains SMN1, SIP1/GEMIN2, DDX20/GEMIN3, GEMIN4, GEMIN5, GEMIN6, GEMIN7, GEMIN8 and STRAP/UNRIP. Interacts with FXR1, FXR2, IGF2BP1, NUFIP1, NUFIP2, MCRS1 and RANBP9. Interacts with CYFIP1 and CYFIP2.,tissue specificity:Highest levels found in neurons, brain, testis, placenta and lymphocytes. Also expressed in epithelial tissues and at very low levels in glial cells.,

Background

The protein encoded by this gene binds RNA and is associated with polysomes. The encoded protein may be involved in mRNA trafficking from the nucleus to the cytoplasm. A trinucleotide repeat (CGG) in the 5' UTR is normally found at 6-53 copies, but an expansion to 55-230 repeats is the cause of fragile X syndrome. Expansion of the trinucleotide repeat may also cause one form of premature ovarian failure (POF1). Multiple alternatively spliced transcript variants that encode different protein isoforms and which are located in different cellular

杭州臻优品生物科技有限公司

热销产品:

蛋白、一抗、抗体对、ELISA试剂盒、生化试剂盒
CCK8试剂盒、QPCR检测试剂盒

检测服务:

ELISA检测及定制服务 | 生化检测 | PCR、QPCR检测 | WB检测
ICO-IP检测 | 切片 | 染色 | 免疫组化 | 免疫荧光 | 透射电镜全套
| 宏基因组、转录组、基因组、蛋白组、代谢组测序


 关注
官网

 关注
客服




locations have been described for this gene. [provided by RefSeq, May 2010],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Mouse testis was stained with anti-FMR1 Rabbit antibody



Human testis was stained with anti-FMR1 Rabbit antibody

杭州臻优品生物科技有限公司

热销产品:

蛋白、一抗、抗体对、ELISA试剂盒、生化试剂盒
CCK8试剂盒、QPCR检测试剂盒

检测服务:


ELISA检测及定制服务 | 生化检测 | PCR、QPCR检测 | WB检测
ICO-IP检测 | 切片 | 染色 | 免疫组化 | 免疫荧光 | 透射电镜全套
| 宏基因组、转录组、基因组、蛋白组、代谢组测序



关注官网



关注客服



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-FMR1 antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: K562 Lane 2: HeLa Predicted band size: 71kDa Observed band size: 70-77kDa

杭州臻优品生物科技有限公司

热销产品:

蛋白、一抗、抗体对、ELISA试剂盒、生化试剂盒
CCK8试剂盒、QPCR检测试剂盒

检测服务:

ELISA检测及定制服务 | 生化检测 | PCR、QPCR检测 | WB检测
ICO-IP检测 | 切片 | 染色 | 免疫组化 | 免疫荧光 | 透射电镜全套
| 宏基因组、转录组、基因组、蛋白组、代谢组测序



关注官网



关注客服