



MLLT6 Rabbit mAb


Catalog No	YP-rAb-17425
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	MLLT6,AF17
Protein Name	Protein AF-17 (ALL1-fused gene from chromosome 17 protein)
Purification Process	Protein A
Specificity	Endogenous
Formulation	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source	Monoclonal, Rabbit,IgG
Dilution	WB 1:1000-1:5000;
Concentration	0.5 mg/ml
Purity	≥90%
Storage Stability	-15° C to -25° C/1 year(Do not lower than -25° C)
Synonyms	AF17
Observed Band	120kD
Calculated Molecular Weight	120kD
Cell Pathway	SUBCELLULAR LOCATION: Nucleus {ECO:0000305}.
Tissue Specificity	
Function	
Background	Enables histone binding activity and nucleosome binding activity. Predicted to be involved in regulation of transcription by RNA polymerase II. Predicted to act upstream of or within several processes, including renal potassium excretion; renal sodium excretion; and renal water absorption. Predicted to be active in nucleus. [provided by Alliance of Genome Resources, Apr 2025]
matters needing attention	Avoid repeated freezing and thawing!





Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Western blot analysis of lysates from NIH/3T3 cell, primary antibody was diluted at 1:1000, 4° over night, Dylight 800 secondary antibody was diluted at 1:10000, 37° 1 hour.

杭州臻优品生物科技有限公司

热销产品:

蛋白、一抗、抗体对、ELISA试剂盒、生化试剂盒
CCK8试剂盒、QPCR检测试剂盒

检测服务:

ELISA检测及定制服务 | 生化检测 | PCR、QPCR检测 | WB检测
ICO-IP检测 | 切片 | 染色 | 免疫组化 | 免疫荧光 | 透射电镜全套
| 宏基因组、转录组、基因组、蛋白组、代谢组测序



关注官网



关注客服