



Methionyl Aminopeptidase 1 Rabbit mAb

Catalog No	YP-rAb-17452
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB,IHC,IF,ELISA
Gene Name	METAP1 KIAA0094
Protein Name	Methionine aminopeptidase 1 (MAP 1) (MetAP 1) (Peptidase M 1)
Purification Process	Protein A
Specificity	Endogenous
Formulation	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source	Monoclonal, Rabbit,IgG
Dilution	IHC 1:200-1:1000; WB 1:2000-1:10000; IF 1:200-1:1000; ELISA 1:5000-1:20000; Note: For IHC, we suggest antigen retrieval with TE buffer pH 9.0
Concentration	0.5 mg/ml
Purity	≥90%
Storage Stability	-15° C to -25° C/1 year(Do not lower than -25° C)
Synonyms	AMPM1 HUMAN,MAP 1,MAP1A,MetAP 1,MetAP1A,Methionine aminopeptidase 1,Peptidase M 1,
Observed Band	42kD
Calculated Molecular Weight	42kD
Cell Pathway	Cytoplasm .
Tissue Specificity	Bone marrow,Placenta,Testis,
Function	Catalytic activity:Release of N-terminal amino acids, preferentially methionine, from peptides and arylamides.,cofactor:Binds 1 sodium ion per subunit. The sodium ion has a structural role.,cofactor:Binds 2 cobalt ions per subunit.,cofactor:Binds 2 cobalt ions per subunit. The true nature of the physiological cofactor is under debate. The enzyme is also active with zinc, manganese or divalent iron ions.,Function:Removes the amino-terminal methionine from nascent proteins.,Function:Removes the amino-terminal methionine from nascent proteins. Required for normal progression through the cell cycle.,similarity:Belongs to the peptidase M24A family.,
Background	catalytic activity:Release of N-terminal amino acids, preferentially methionine, from peptides and arylamides.,cofactor:Binds 1 sodium ion per subunit. The sodium ion has a structural role.,cofactor:Binds 2 cobalt ions per

杭州臻优品生物科技有限公司

热销产品:

蛋白、一抗、抗体对、ELISA试剂盒、生化试剂盒
CCK8试剂盒、QPCR检测试剂盒

检测服务:

ELISA检测及定制服务 | 生化检测 | PCR、QPCR检测 | WB检测
| CO-IP检测 | 切片 | 染色 | 免疫组化 | 免疫荧光 | 透射电镜全套
| 宏基因组、转录组、基因组、蛋白组、代谢组测序



关注官网



关注客服



subunit.,cofactor: Binds 2 cobalt ions per subunit. The true nature of the physiological cofactor is under debate. The enzyme is also active with zinc, manganese or divalent iron ions.,function: Removes the amino-terminal methionine from nascent proteins.,function: Removes the amino-terminal methionine from nascent proteins. Required for normal progression through the cell cycle.,similarity: Belongs to the peptidase M24A family.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Western blot analysis of lysates from A375 cell, primary antibody was diluted at 1:1000, 4° over night, Dylight 800 secondary antibody was diluted at 1:10000, 37° 1hour.

