



C3orf38 Rabbit mAb

Catalog No	YP-rAb-17662
Isotype	IgG
Reactivity	Human
Applications	WB
Gene Name	C3orf38
Protein Name	Uncharacterized protein C3orf38
Purification Process	Protein A
Specificity	Endogenous
Formulation	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source	Monoclonal, Rabbit,IgG
Dilution	WB 1:1000-5000
Concentration	0.5 mg/ml
Purity	≥90%
Storage Stability	-15° C to -25° C/1 year(Do not lower than -25° C)
Synonyms	C3orf38 ; Uncharacterized protein C3orf38 ;
Observed Band	36kD
Calculated Molecular Weight	
Cell Pathway	
Tissue Specificity	
Function	May be involved in apoptosis regulation. {ECO:0000269 PubMed:17464193}.
Background	Involved in positive regulation of apoptotic process. Located in nucleus. [provided by Alliance of Genome Resources, Feb 2025]
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

杭州臻优品生物科技有限公司

热销产品:

蛋白、一抗、抗体对、ELISA试剂盒、生化试剂盒
CCK8试剂盒、QPCR检测试剂盒

检测服务:


ELISA检测及定制服务 | 生化检测 | PCR、QPCR检测 | WB检测
ICO-IP检测 | 切片 | 染色 | 免疫组化 | 免疫荧光 | 透射电镜全套
| 宏基因组、转录组、基因组、蛋白组、代谢组测序



关注官网



关注客服



Western Blot analysis of L02 whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-C3orf38 rabbit mAb. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody.

