



BTNL3 Rabbit mAb


Catalog No	YP-rAb-17675
Isotype	IgG
Reactivity	Human
Applications	WB,FC
Gene Name	BTNL3;BTNLR;COLF4100;UNQ744/PRO1472
Protein Name	Butyrophilin-like protein 3;Butyrophilin-like receptor;
Purification Process	Protein A
Specificity	Endogenous
Formulation	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source	Monoclonal, Rabbit,IgG
Dilution	WB 1:1000-5000; FC 1:100-300
Concentration	0.5 mg/ml
Purity	≥90%
Storage Stability	-15° C to -25° C/1 year(Do not lower than -25° C)
Synonyms	BTNL3 ; BTNLR ; COLF4100 ; UNQ744/PRO1472 ; Butyrophilin-like protein 3 ; Butyrophilin-like receptor ;
Observed Band	51kD
Calculated Molecular Weight	
Cell Pathway	Membrane; Single-pass type I membrane protein.
Tissue Specificity	Expressed in small intestine, colon, testis, spleen, and leukocyte. {ECO:0000269 PubMed:10429365}.
Function	
Background	Predicted to enable signaling receptor binding activity. Predicted to be involved in T cell receptor signaling pathway and regulation of cytokine production. Predicted to be located in membrane. Predicted to be active in external side of plasma membrane. [provided by Alliance of Genome Resources, Feb 2025]
matters needing attention	Avoid repeated freezing and thawing!





Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Western Blot analysis of HeLa whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-BTNL3 rabbit mAb. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody.

杭州臻优品生物科技有限公司

热销产品:

蛋白、一抗、抗体对、ELISA试剂盒、生化试剂盒
CCK8试剂盒、QPCR检测试剂盒

检测服务:

ELISA检测及定制服务 | 生化检测 | PCR、QPCR检测 | WB检测
ICO-IP检测 | 切片 | 染色 | 免疫组化 | 免疫荧光 | 透射电镜全套
| 宏基因组、转录组、基因组、蛋白组、代谢组测序



关注官网



关注客服



杭州臻优品生物科技有限公司

热销产品:

蛋白、一抗、抗体对、ELISA试剂盒、生化试剂盒
CCK8试剂盒、QPCR检测试剂盒

检测服务:

ELISA检测及定制服务 | 生化检测 | PCR、QPCR检测 | WB检测
ICO-IP检测 | 切片 | 染色 | 免疫组化 | 免疫荧光 | 透射电镜全套
| 宏基因组、转录组、基因组、蛋白组、代谢组测序



关注官网



关注客服